Index of Authors

AHKONG, Q. F. *see* QUIRK, S. J. 159–167

AHMED, K., WILSON, M. J., GOUELI, S. A. & WILLIAMS-ASHMAN, H. G. Effects of polyamines on prostatic chromatin- and non-histone-protein-associated protein kinase reactions 739–750

ALI, S. Y. *see* BAYLISS, M. T. 683–693

ANDERSSON, A. Opposite effects of starvation on oxidation of [^{14}C]adenosine and adenosine-induced insulin release by isolated mouse pancreatic islets 619–621

ASHBY, P., BENNETT, D. P., SPENCER, I. M. & ROBINSON, D. S. Post-translational regulation of lipoprotein lipase activity in adipose tissue 865–872

ASINA, S. *see* JAIN, K. 31–37

AUSTIN, L. *see* KWOK, C. T. 15–22

AZOULAY, E. *see* GIORDANO, G. 553–561

BALLARD, F. J. *see* CHANDLER, C. S. 151–158

BARRITT, G. J. *see* HUGHES, B. P. 295–304

BARTFAI, T., BREAKEFIELD, X. O. & GREENGARD, P. Regulation of synthesis of guanosine 3′:5′-cyclic monophosphate in neuroblastoma cells 119–127

BATES, D. J., PERRETT, D. & MOWBRAY, J. Systematic variations in the content of the purine nucleotides in the steady-state perfused rat heart. Evidence for the existence of controlled storage and release of adenine nucleotides 485–493

BATES, P. C. *see* LAURENT, G. J. 393–401, 419–427

BAYLISS, M. T. & ALI, S. Y. Age-related changes in the composition and structure of human articular-cartilage proteoglycans 683–693

BEEBEE, T. J. C. The use of rat liver nucleoplasm for the characterization of heterogeneous nuclear ribonucleic acid synthesis *in vitro* 715–725

BENITO, M. & WILLIAMSON, D. H. Evidence for a reciprocal relationship between lipogenesis and ketogenesis in hepatocytes from fed virgin and lactating rats 331–334

BENNETT, D. P. *see* ASHBY, P. 865–872

BERTHELIER, M. *see* THOMOPOULOS, P. 169–174

BETTERIDGE, A. & WALLIS, M. Stimulation of anterior pituitary prostaglandin E content and somatotropin (growth hormone) synthesis by phospholipase A 319–323

BING, R. F. & SCHULSTER, D. Adenosine 3′:5′-cyclic monophosphate production and steroidogenesis by isolated rat adrenal glomerulosa cells. Effects of angiotensin II and [Sar^1,Ala^8]angiotensin II 39–45

BLACKMORE, P. F. *see* WILLIAMS, J. F. 241–256, 257–282

BLACKWELL, C. M. & TURNER, J. M. Microbial metabolism of amino alcohols. Formation of coenzyme B_{12}-dependent ethanolamine ammonia-lyase and its concerted induction in *Escherichia coli* 751–757

BONNEY, R. J., WIGHTMAN, P. D., DAVIES, P., SADOWSKI, S. J., KUEHL, F. A., JR. & HUMES, J. L. Regulation of prostaglandin synthesis and of the selective release of lysosomal hydrolases by mouse peritoneal macrophages 433–442

BOOTH, A. *see* MCMILLAN, R. M. 595–598

BOOTH, R. F. G. & CLARK, J. B. A rapid method for the preparation of relatively pure metabolically competent synaptosomes from rat brain 365–370

BORUM, P. R. Variation in tissue carnitine concentrations with age and sex in the rat 677–681

BOSCHERO, A. C. *see* SENER, A. 217–232

BOTHAM, G. M. *see* QUIRK, S. J. 159–167

BOURBON, J. *see* GILBERT, M. 785–789

BRADFORD, N. M. *see* JOSEPH, S. K. 827–836

BREAKEFIELD, X. O. *see* BARTFAI, T. 119–127; CROOKS, P. A. 187–196

BRINDLEY, D. N. *see* GLENNY, H. P. 777–784

BROWN, S. B. & THOMAS, S. E. The mechanism of haem degradation *in vitro*. Kinetic evidence for the formation of a haem–oxygen complex 327–330

BRYŁA, J. *see* ZALESKI, J. 563–568

BUCKLEY, W. T. & MILLIGAN, L. P. Participation of cysteine and cystine in inactivation of tyrosine aminotransferase in rat liver homogenates 449–454

BUCKSHEE, K. *see* KASID, A. 531–539

BUDDECKE, E. *see* PRINZ, R. 671–676

BURGESS, R. J., WALKER, J. H. & MAYER, R. J. Choice of precursors for the measurement of protein turnover by the double-isotope method. Application to the study of mitochondrial proteins 919–926

BURGESS, R. J. *see also* WALKER, J. H. 927–932

BURNSIDE, J. *see* SCHNEIDER, D. L. 75–82

BYGRAVE, F. L. *see* PRPIĆ, V. 705–714

CAMPBELL, A. K. & DORMER, R. L. Inhibition by calcium ions of adenosine cyclic monophosphate formation in sealed pigeon erythrocyte 'ghosts'. A study using the photoprotein obelin 53–66

CARTENÌ-FARINA, M. *see* DE ROSA, M. 1–7

CASTIGLIONE, C. M. *see* CROOKS, P. A. 187–196

CHANDLER, C. S. & BALLARD, F. J. Accelerated breakdown of reticulocyte protein formed under conditions of amino acid depletion 151–158

CHAPMAN, M. J. *see* THOMOPOULOS, P. 169–174

CHATTOPADHYAY, S. *see* SAMANTA, T. B. 593–594

CHAUDHURI, G. *see* MYATT, L. 523–529

CHIN, J. G. *see* DENTON, R. M. 899–906

CLARK, J. B. *see* BOOTH, R. F. G. 365–370; PATEL, T. B. 951–958

CLARK, M. G. & JARRETT, I. G. Responsiveness to glucagon by isolated rat hepatocytes controlled by the redox state of the cytosolic nicotinamide–adenine dinucleotide couple acting on adenosine 3′:5′-cyclic monophosphate phosphodiesterase 805–816

CLARK, M. G. *see also* WILLIAMS, J. F. 241–256, 257–282

CLIFFORD, A. J. *see* HOOVER-PLOW, J. L. 137–142

COIA, A. A. *see* LEADER, D. P. 569–572

COWARD, J. K. *see* CROOKS, P. A. 187–196

CROOK, R. B. & TOMKINS, G. M. Effect of glutamine on the degradation of glutamine synthetase in hepatoma tissue-culture cells 47–52

CROOKS, P. A., BREAKEFIELD, X. O., SULENS, C. H., CASTIGLIONE, C. M. & COWARD, J. K. Extensive conjugation of dopamine (3,4-dihydroxyphenethylamine) metabolites in cultured human skin fibroblasts and rat hepatoma cells 187–196

CURTIS, C. G. *see* DESAI, R. 455–462; POWELL, G. M. 443–448

DAS, C. *see* RAMAKRISHNAN, S. 599–602
DAVIDSON, M. M. L. *see* HASLAM, R. J. 83–95
DAVIES, P. *see* BONNEY, R. J. 433–442
DAWSON, R. M. C. *see* IRVINE, R. F. 475–484
DAZORD, A., GALLET, D. & SAEZ, J. M. Effect of corticotropin treatment *in vivo* on the synthesis of a specific adrenal cytosolic protein. Characterization by dual-labelling technique and polyacrylamide-gel electrophoresis 233–239
DE ROSA, M., DE ROSA, S., GAMBACORTA, A., CARTENÌ-FARINA, M. & ZAPPIA, V. The biosynthetic pathway of new polyamines in *Caldariella acidophila* 1–7
DE ROSA, S. *see* DE ROSA, M. 1–7
DECHATELET, L. R. *see* IVERSON, D. B. 175–178
DENARO, M. *see* MONTANARO, L. 371–379
DENTON, R. M., RICHARDS, D. A. & CHIN, J. G. Calcium ions and the regulation of NAD^+-linked isocitrate dehydrogenase from the mitochondria of rat heart and other tissues 899–906
DENTON, R. M. *see also* HEMS, D. A. 627–629
DESAI, R., TETLEY, T. D., CURTIS, C. G., POWELL, G. M. & RICHARDS, R. J. Studies on the fate of pulmonary surfactant in the lung 455–462
DESCHRYVER, C. *see* SCHLESINGER, P. H. 103–109
DESJARDINS, J. V. *see* HASLAM, R. J. 83–95
DEVIS, G. *see* SENER, A. 217–232
DODD, G. *see* MENEVSE, A. 845–854
DOEBBER, T. W. *see* SCHLESINGER, P. H. 103–109
DORMER, R. L. *see* CAMPBELL, A. K. 53–66
DOU, J. H. *see* GIORDANO, G. 553–561
DUNHAM, S. M. & THURSTON, C. F. Control of isocitrate lyase synthesis in *Chlorella fusca* var. *vacuolata*. Rate of enzyme synthesis in the presence and absence of acetate measured by [^{35}S]methionine labelling and immunoprecipitation 179–185

EDELSTEIN, S., NOFF, D., FREEMAN, D., SHEVES, M. & MAZUR, Y. Synthesis of 1α-hydroxy[7-^{3}H]cholecalciferol and its metabolism in the chick 111–117
ELDER, M. G. *see* MYATT, L. 523–529
ELLAR, D. J. *see* HOGARTH, C. 197–203
ELLIOTT, K. R. F. *see* SMITH, S. A. 817–825
ELLIOTT, W. H. *see* PEARCE, P. D. 611–614

FELÍU, J. E. *see* HUE, L. 791–797
FERRÉ, P., PÉGORIER, J.-P., WILLIAMSON, D. H. & GIRARD, J. R. The development of ketogenesis at birth in the rat 759–765
FERRÉ, P. & WILLIAMSON, D. H. Evidence for the participation of aspartate aminotransferase in hepatic glucose synthesis in the suckling newborn rat 335–338
FEY, S. *see* HUVOS, P. 381–391
FISCHER, J. & SHARP, G. W. G. Effects of cholera toxin and guanosine 5′-[βγ-imido]triphosphate on β-adrenergic-receptor affinity 505–510
FREEMAN, D. *see* EDELSTEIN, S. 111–117
FUNAE, Y. *see* FUNAKAWA, S. 977–981
FUNAKAWA, S., FUNAE, Y. & YAMAMOTO, K. Conversion between renin and high-molecular-weight renin in the dog 977–981

GALLAGHER, K. *see* TENG, C. S. 1003–1006
GALLET, D. *see* DAZORD, A. 233–239
GAMBACORTA, A. *see* DE ROSA, M. 1–7
GARLICK, P. J. An analysis of errors in estimation of the rate of protein synthesis by constant infusion of a labelled amino acid 402–405
GARTON, G. A. *see* SCAIFE, J. R. 799–804
GASPERI-CAMPANI, A. *see* MONTANARO, L. 371–379
GILBERT, M. & BOURDON, J. Turnover of liver glycogen in the rat foetus 785–789
GIORDANO, G., GRILLET, L., ROSSET, R., DOU, J. H., AZOULAY, E. & HADDOCK, B. A. Characterization of an *Escherichia coli* K12 mutant that is sensitive to chlorate when grown aerobically 553–561
GIRARD, J. R. *see* FERRÉ, P. 759–765; ROBINSON, A. M. 343–346
GIUDICELLI, Y. Thyroid-hormone modulation of the number of β-adrenergic receptors in rat fat-cell membranes 1007–1010
GLENNY, H. P. & BRINDLEY, D. N. The effects of cortisol, corticotropin and thyroxine on the synthesis of glycerolipids and on the phosphatidate phosphohydrolase activity in rat liver 777–784
GLINSMANN, W. H. *see* TAO, S.-H. 347–350
GOLDSPINK, D. F. The effects of food deprivation on protein turnover and nucleic acid concentrations of active and immobilized extensor digitorum longus muscles of the rat 603–606
GOMEZ-FERNANDEZ, J. C. & HARRIS, D. A. A thermodynamic analysis of the interaction between the mitochondrial coupling adenosine triphosphatase and its naturally occurring inhibitor protein 967–975
GORDON, J. L. *see* MCMILLAN, R. M. 595–598
GORGA, F. R. *see* SCHNEIDER, D. L. 75–82
GOUELI, S. A. *see* AHMED, K. 739–750
GRANDCHAMP, B., PHUNG, N. & NORDMANN, Y. The mitochondrial localization of coproporphyrinogen III oxidase 97–102
GRANT, M. E. *see* HEATHCOTE, J. G. 283–294
GREENGARD, P. *see* BARTFAI, T. 119–127
GRIFFIN, H. D. & HAWTHORNE, J. N. Calcium-activated hydrolysis of phosphatidyl-*myo*-inositol 4-phosphate and phosphatidyl-*myo*-inositol 4,5-bisphosphate in guinea-pig synaptosomes 541–552
GRILLET, L. *see* GIORDANO, G. 553–561

HADDOCK, B. A. *see* GIORDANO, G. 553–561
HALES, C. N. *see* MCPHERSON, M. A. 855–863
HALL, C. & LIM, L. The metabolism of high-molecular-weight ribonucleic acid in hypothalamic and cortical regions of the developing female rat brain 511–521
HALLIDAY, D. *see* KREBS, H. A. 733–737
HARDWICKE, P. M. D. *see* HUVOS, P. 381–391
HARRIS, D. A. *see* GOMEZ-FERNANDEZ, J. C. 967–975
HARRIS, E. J. Anion/calcium ion ratios and proton production in some mitochondrial calcium ion uptakes 983–991
HASLAM, R. J., DAVIDSON, M. M. L. & DESJARDINS, J. V. Inhibition of adenylate cyclase by adenosine analogues in preparations of broken and intact human platelets. Evidence for the unidirectional control of platelet function by cyclic AMP 83–95
HASSINEN, I. E. *see* SAVOLAINEN, M. J. 885–892
HAWTHORNE, J. N. *see* GRIFFIN, H. D. 541–552

HEATHCOTE, J. G., SEAR, C. H. J. & GRANT, M. E. Studies on the assembly of the rat lens capsule. Biosynthesis and partial characterization of the collagenous components 283–294
HEMINGTON, N. *see* IRVINE, R. F. 475–484
HEMS, D. A., MCCORMACK, J. G. & DENTON, R. M. Activation of pyruvate dehydrogenase in the perfused rat liver by vasopressin 627–629
HEMS, D. A. *see also* WHITTON, P. D. 893–898
HEMS, R. *see* KREBS, H. A. 733–737
HERCHUELZ, A. *see* SENER, A. 217–232
HERS, H.-G. *see* HUE, L. 791–797
HILF, R. *see* HISSIN, P. J. 205–215
HINGORANI, V. *see* KASID, A. 531–539
HISSIN, P. J. & HILF, R. α-Aminoisobutyrate transport into cells from R3230AC mammary adenocarcinoma. Evidence for sodium ion-dependent and -independent carrier-mediated entry and effects of diabetes 205–215
HOGARTH, C. & ELLAR, D. J. Calcium accumulation during sporulation of *Bacillus megaterium* KM 197–203
HOOVER-PLOW, J. L. & CLIFFORD, A. J. The effect of surgical trauma on muscle protein turnover in rats 137–142
HOUSTON, B. *see* NIMMO, H. G. 607–610
HOZUMI, M. *see* TOMIDA, M. 665–669
HUANG, F. L. *see* TAO, S.-H. 347–350
HUANG, M.-T. *see* MACDONALD, M. J. 495–504
HUE, L., FELÍU, J. E. & HERS, H.-G. Control of gluconeogenesis and of enzymes of glycogen metabolism in isolated rat hepatocytes. A parallel study of the effect of phenylephrine and of glucagon 791–797
HUGHES, B. P. & BARRITT, G. J. Effects of glucagon and $N^6O^{2'}$-dibutyryladenosine 3′:5′-cyclic monophosphate on calcium transport in isolated rat liver mitochondria 295–304
HUGHES, J. & MELLOWS, G. Inhibition of isoleucyl-transfer ribonucleic acid synthetase in *Escherichia coli* by pseudomonic acid 305–318
HUMES, J. L. *see* BONNEY, R. J. 433–442
HUTTON, J. C. *see* SENER, A. 217–232
HUVOS, P., FEY, S. & HARDWICKE, P. M. D. Effect of various methods of preparation on the apparent protein composition of eukaryotic ribosomes. An essential preliminary to stoicheiometric measurements 381–391

INFANTE, J. P. & KINSELLA, J. E. Control of phosphatidylcholine synthesis and the regulatory role of choline kinase in rat liver. Evidence from essential-fatty acid-deficient rats 631–633
IRVINE, R. F., HEMINGTON, N. & DAWSON, R. M. C. The hydrolysis of phosphatidylinositol by lysosomal enzymes of rat liver and brain 475–484
IVERSON, D. B., WANG-IVERSON, P., SPITZNAGEL, J. K. & DECHATELET, L. R. Subcellular localization of NAD(P)H oxidase(s) in human neutrophilic polymorphonuclear leucocytes 175–178

JAIN, K., ASINA, S. & LOGOTHETOPOULOS, J. Stimulation of proinsulin biosynthesis and insulin release by pyruvate and lactate 31–37
JARRETT, I. G. *see* CLARK, M. G. 805–816
JOCELYN, P. C. The reduction of diamide by rat liver mitochondria and the role of glutathione 649–664
JOHNSON, T. C. *see* MILENKOVIC, A. G. 695–704
JOSEPH, S. K., BRADFORD, N. M. & MCGIVAN, J. D. Characteristics of the transport of alanine, serine and glutamine across the plasma membrane of isolated rat liver cells 827–836
JOSEPH, S. K. & MCGIVAN, J. D. The effects of ammonium chloride and bicarbonate on the activity of glutaminase in isolated liver mitochondria 837–844

KANAGASUNTHERAM, P. & LIM, S. C. Calcium-dependent inhibition of protein synthesis in rat parotid gland 23–29
KASID, A., BUCKSHEE, K., HINGORANI, V. & LAUMAS, K. R. Interaction of progestins with steroid receptors in human uterus 531–539
KAWAZU, S. *see* SENER, A. 217–232
KEIR, H. M. *see* MELVIN, W. T. 933–941
KEULEMANS, K. *see* MULDER, G. J. 959–965
KINSELLA, J. E. *see* INFANTE, J. P. 631–633
KREBS, H. A., HEMS, R., LUND, P., HALLIDAY, D. & READ, W. W. C. Sources of ammonia for mammalian urea synthesis 733–737
KUEHL, F. A., JR. *see* BONNEY, R. J. 433–442
KWOK, C. T. & AUSTIN, L. Phospholipid composition and metabolism in mouse muscular dystrophy 15–22

LAGRANGE, D. *see* THOMOPOULOS, P. 169–174
LARDY, H. A. *see* MACDONALD, M. J. 495–504
LATIF, A. *see* NAKHLA, A. M. 339–342
LAUDAT, M.-H. *see* THOMOPOULOS, P. 169–174
LAUMAS, K. R. *see* KASID, A. 531–539
LAURENT, G. J., SPARROW, M. P., BATES, P. C. & MILLWARD, D. J. Turnover of muscle protein in the fowl. Collagen content and turnover in cardiac and skeletal muscles of the adult fowl and the changes during stretch-induced growth 419–427
LAURENT, G. J., SPARROW, M. P., BATES, P. C. & MILLWARD, D. J. Turnover of muscle protein in the fowl (*Gallus domesticus*). Rates of protein synthesis in fast and slow skeletal, cardiac and smooth muscle of the adult fowl 393–401
LAURENT, G. J., SPARROW, M. P. & MILLWARD, D. J. Turnover of muscle protein in the fowl. Changes in rates of protein synthesis and breakdown during hypertrophy of the anterior and posterior latissimus dorsi muscles 407–417
LEADER, D. P. & COIA, A. A. The acidic ribosomal phosphoprotein of eukaryotes and its relationship to ribosomal proteins L7 and L12 of *Escherichia coli* 569–572
LEHTOVAARA, P. Oxidation of glycine by *Phaseolus* leghaemoglobin with associated catabolic reactions at the haem 351–358
LEHTOVAARA, P. & PERTTILÄ, U. Bile-pigment formation from different leghaemoglobins. Methine-bridge specificity of coupled oxidation 359–364
LIM, L. *see* HALL, C. 511–521; MYATT, L. 523–529
LIM, S. C. *see* KANAGASUNTHERAM, P. 23–29
LOGOTHETOPOULOS, J. *see* JAIN, K. 31–37
LOPEZ, A. *see* PERRY, B. N. 873–883
LUCY, J. A. *see* QUIRK, S. J. 159–167
LUND, P. *see* KREBS, H. A. 733–737
LYNCH, A. *see* TAO, S.-H. 347–350

MacDonald, M. J., Huang, M.-T. & Lardy, H. A. Hyperglycaemic activity and metabolic effects of 3-aminopicolinic acid 495–504
MacIntyre, D. E. *see* McMillan, R. M. 595–598
Malaisse, W. J. *see* Sener, A. 217–232
Mandell, B. F. *see* Schlesinger, P. H. 103–109
Maruyama, K., Sunde, M. L. & Swick, R. W. Growth and muscle protein turnover in the chick 573–582
May, B. K. *see* Pearce, P. D. 611–614
Mayer, R. J. *see* Burgess, R. J. 919–926; Walker, J. H. 927–932
Mazur, Y. *see* Edelstein, S. 111–117
McCormack, J. G. *see* Hems, D. A. 627–629
McGivan, J. D. *see* Joseph, S. K. 827–836, 837–844
McMillan, R. M., MacIntyre, D. E., Booth, A. & Gordon, J. L. Malonaldehyde formation in intact platelets is catalysed by thromboxane synthase 595–598
McPherson, M. A. & Hales, C. N. Control of amylase biosynthesis and release in the parotid gland of the rat 855–863
Mellows, G. *see* Hughes, J. 305–318
Melvin, W. T. & Keir, H. M. Onset of ribosome degradation during cessation of growth in BHK-21/C13 cells 933–941
Menevse, A., Dodd, G. & Poynder, T. M. A chemical-modification approach to the olfactory code. Studies with a thiol-specific reagent 845–854
Milenkovic, A. G., Rachmeler, M. & Johnson, T. C. Growth and metabolism of fucosylated plasma-membrane glycoproteins in mouse neuroblastoma N2a cells 695–704
Miller, M. J. *see* Schlesinger, P. H. 103–109
Milligan, L. P. *see* Buckley, W. T. 449–454
Millward, D. J. *see* Laurent, G. J. 393–401, 407–417, 419–427
Montanaro, L., Sperti, S., Zamboni, M., Denaro, M., Testoni, G., Gasperi-Campani, A. & Stirpe, F. Effect of modeccin on the steps of peptide-chain elongation 371–379
Mowbray, J. *see* Bates, D. J. 485–493
Mulder, G. J. & Keulemans, K. Metabolism of inorganic sulphate in the isolated perfused rat liver. Effect of sulphate concentration on the rate of sulphation by phenol sulphotransferase 959–965
Murray, A. W. & Rogers, A. Calcium-dependent protein modulator of cyclic nucleotide phosphodiesterases from mouse epidermis 727–732
Myatt, L., Chaudhuri, G., Elder, M. G. & Lim, L. The oestrogen receptor in the rat uterus in relation to intra-uterine devices and the oestrous cycle 523–529

Nakhla, A. M. & Latif, A. A possible role for 5-hydroxytryptamine as a mediator for calcitonin actions on the gastrointestinal tract and pancreas in rats 339–342
Nettleton, C. J. *see* Schneider, D. L. 75–82
Newsholme, E. A. & Williams, T. The role of phosphoenolpyruvate carboxykinase in amino acid metabolism in muscle 623–626
Nicholas, R. A. & Ordal, G. W. Inhibition of bacterial transport by uncouplers of oxidative phosphorylation. Effects of pentachlorophenol and analogues in *Bacillus subtilis* 639–647
Nicholls, D. G. The regulation of extramitochondrial free calcium ion concentration by rat liver mitochondria 463–474
Nicholls, D. M. *see* Petryshyn, R. A. 907–917
Nimmo, H. G. & Houston, B. Rat adipose-tissue glycerol phosphate acyltransferase can be inactivated by cyclic AMP-dependent protein kinase 607–610
Noff, D. *see* Edelstein, S. 111–117
Nordmann, Y. *see* Grandchamp, B. 97–102

Olavesen, A. H. *see* Powell, G. M. 443–448
Ord, M. G. & Stocken, L. A. Initiation of transcription in normal and phosphorylated rat liver nucleosomes 615–618
Ordal, G. W. *see* Nicholas, R. A. 639–647

Palmer, D. N., Rabin, B. R. & Williams, D. J. A subpopulation of rat liver membrane-bound ribosomes that are detached *in vitro* by carcinogens and centrifugation 9–14
Patel, T. B. & Clark, J. B. Acetoacetate metabolism in rat brain. Development of acetoacetyl-coenzyme A deacylase and 3-hydroxy-3-methylglutaryl-coenzyme A synthase 951–958
Pearce, P. D., May, B. K. & Elliott, W. H. Proteinase-sensitive release of enzymes from pancreatic microsomal fraction 611–614
Pégorier, J.-P. *see* Ferré, P. 759–675
Perrett, D. *see* Bates, D. J. 485–493
Perry, B. N. & Lopez, A. The binding of ^{3}H-labelled oestradiol- and progesterone-receptor complexes to hypothalamic chromatin of male and female sheep 873–883
Perttilä, U. *see* Lehtovaara, P. 359–364
Petryshyn, R. A. & Nicholls, D. M. Studies of a factor from dystrophic mouse muscle inhibitory towards protein synthesis 907–917
Phung, N. *see* Grandchamp, B. 97–102
Pogson, C. I. *see* Smith, S. A. 817–825
Powell, G. M., Olavesen, A. H. & Curtis, C. G. Biliary excretion of cyclohexylphenyl 4-[^{35}S]sulphate in the guinea pig 443–448
Powell, G. M. *see also* Desai, R. 455–462
Poynder, T. M. *see* Menevse, A. 845–854
Prebble, J. *see* Ramadan-Talib, Z. 767–775
Prinz, R., Schwermann, J., Buddecke, E. & von Figura, K. Endocytosis of sulphated proteoglycans by cultured skin fibroblasts 671–676
Prpić, V., Spencer, T. L. & Bygrave, F. L. Stable enhancement of calcium retention in mitochondria isolated from rat liver after the administration of glucagon to the intact animal 705–714

Quirk, S. J., Ahkong, Q. F., Botham, G. M., Vos, J. & Lucy, J. A. Membrane proteins in human erythrocytes during cell fusion induced by oleoylglycerol 159–167

Rabin, B. R. *see* Palmer, D. N. 9–14
Rachmeler, M. *see* Milenkovic, A. G. 695–704
Ramadan-Talib, Z. & Prebble, J. Photosensitivity of respiration in *Neurospora* mitochondria. A protective role for carotenoid 767–775

RAMAKRISHNAN, S., DAS, C. & TALWAR, G. P. Recognition of the beta-subunit of human chorionic gonadotropin and sub-determinants by target tissue receptors 599–602
READ, W. W. C. *see* KREBS, H. A. 733–737
REID, R. A. *see* SMITH, G. S. 1011–1014
RICHARDS, D. A. *see* DENTON, R. M. 899–906
RICHARDS, R. J. *see* DESAI, R. 455–462
ROBINSON, A. M., GIRARD, J. R. & WILLIAMSON, D. H. Evidence for a role of insulin in the regulation of lipogenesis in lactating rat mammary gland. Measurements of lipogenesis *in vivo* and plasma hormone concentrations in response to starvation and refeeding 343–346
ROBINSON, A. M. & WILLIAMSON, D. H. Utilization of D-3-hydroxy[3-^{14}C]butyrate for lipogenesis *in vivo* in lactating rat mammary gland 635–638
ROBINSON, D. S. *see* ASHBY, P. 865–872
RODMAN, J. S. *see* SCHLESINGER, P. H. 103–109
RODRIGUES, L. M. *see* WHITTON, P. D. 893–898
ROGERS, A. *see* MURRAY, A. W. 727–732
ROSSET, R. *see* GIORDANO, G. 553–561
ROY, N. *see* SAMANTA, T. B. 593–594

SADOWSKI, S. J. *see* BONNEY, R. J. 433–442
SAEZ, J. M. *see* DAZORD, A. 233–239
SAMANTA, T. B., ROY, N. & CHATTOPADHYAY, S. An improved 11α-hydroxylation of progesterone by *Aspergillus ochraceus* TS 593–594
SAVOLAINEN, M. J. & HASSINEN, I. E. Mechanisms for the effects of ethanol on hepatic phosphatidate phosphohydrolase 885–892
SCAIFE, J. R., WAHLE, K. W. J. & GARTON, G. A. Utilization of methylmalonate for the synthesis of branched-chain fatty acids by preparations of chicken liver and sheep adipose tissue 799–804
SCHLESINGER, P. H., DOEBBER, T. W., MANDELL, B. F., WHITE, R., DESCHRYVER, C., RODMAN, J. S., MILLER, M. J. & STAHL, P. Plasma clearance of glycoproteins with terminal mannose and *N*-acetylglucosamine by liver non-parenchymal cells. Studies with β-glucuronidase, *N*-acetyl-β-D-glucosaminidase, ribonuclease B and agalacto-orosomucoid 103–109
SCHNEIDER, D. L., BURNSIDE, J., GORGA, F. R. & NETTLETON, C. J. Properties of the membrane proteins of rat liver lysosomes. The majority of lysosomal membrane proteins are exposed to the cytoplasm 75–82
SCHULSTER, D. *see* BING, R. F. 39–45
SCHWERMANN, J. *see* PRINZ, R. 671–676
SEAR, C. H. J. *see* HEATHCOTE, J. G. 283–294
SENER, A., KAWAZU, S., HUTTON, J. C., BOSCHERO, A. C., DEVIS, G., SOMERS, G., HERCHUELZ, A. & MALAISSE, W. J. The stimulus–secretion coupling of glucose-induced insulin release. Effect of exogenous pyruvate on islet function 217–232
SHARP, G. W. G. *see* FISCHER, J. 505–510
SHEVES, M. *see* EDELSTEIN, S. 111–117
SMITH, G. S. & REID, R. A. The influence of respiratory state on monoamine oxidase activity in rat liver mitochondria 1011–1014
SMITH, S. A., ELLIOTT, K. R. F. & POGSON, C. I. Differential effects of tryptophan on glucose synthesis in rats and guinea pigs 817–825
SOMERS, G. *see* SENER, A. 217–232
SPARROW, M. P. *see* LAURENT, G. J. 393–401, 407–417, 419–427
SPEAKE, B. K. & WHITE, D. A. Lipid-linked oligosaccharides containing glucose in lactating rabbit mammary gland 993–1000
SPENCER, I. M. *see* ASHBY, P. 865–872
SPENCER, T. L. *see* PRPIĆ, V. 705–714
SPERTI, S. *see* MONTANARO, L. 371–379
SPITZNAGEL, J. K. *see* IVERSON, D. B. 175–178
STAHL, P. *see* SCHLESINGER, P. H. 103–109
STIRPE, F. *see* MONTANARO, L. 371–379
STOCKEN, L. A. *see* ORD, M. G. 615–618
STRANGE, P. G. A method for measuring relative changes in guanosine 3′:5′-cyclic monophosphate in mouse neuroblastoma cells on muscarinic cholinergic stimulation 583–590
SULENS, C. H. *see* CROOKS, P. A. 187–196
SUNDE, M. L. *see* MARUYAMA, K. 573–582
SWICK, R. W. *see* MARUYAMA, K. 573–582

TAKENAGA, K. *see* TOMIDA, M. 665–669
TALWAR, G. P. *see* RAMAKRISHNAN, S. 599–602
TAO, S.-H., HUANG, F. L., LYNCH, A. & GLINSMANN, W. H. Control of rat skeletal-muscle phosphorylase phosphatase activity by adrenaline 347–350
TAYLOR, W. H. *see* WALKER, V. 429–432
TENG, C. S., GALLAGHER, K. & TENG, C. T. Isolation of a high-molecular-weight high-mobility-group-type non-histone protein from hen oviduct 1003–1006
TENG, C. S. & TENG, C. T. Studies on sex-organ development. Oestrogenic effect on ornithine decarboxylase activity in the differentiating Müllerian ducts and other organs of the chick embryo 143–149
TENG, C. T. *see* TENG, C. S. 143–149, 1003–1006
TESTONI, G. *see* MONTANARO, L. 371–379
TETLEY, T. D. *see* DESAI, R. 455–462
THOMAS, S. E. *see* BROWN, S. B. 327–330
THOMOPOULOS, P., BERTHELIER, M., LAGRANGE, D., CHAPMAN, M. J. & LAUDAT, M.-H. Stimulation of triacylglycerol synthesis in rat adipocytes by plasma very-low-density lipoproteins 169–174
THURSTON, C. F. *see* DUNHAM, S. M. 179–185
TOMIDA, M., TAKENAGA, K., YAMAMOTO, Y. & HOZUMI, M. Enhancement by double-stranded polyribonucleotides of production by cultured mouse peritoneal macrophages of differentiation-stimulating factor(s) for mouse myeloid leukaemic cells 665–669
TOMKINS, G. M. *see* CROOK, R. B. 47–52
TUAN, R. S. & ZRIKE, J. Functional involvement of carbonic anhydrase in calcium transport of the chick chorioallantoic membrane 67–74
TURNER, J. M. *see* BLACKWELL, C. M. 751–757

VON FIGURA, K. & WEBER, E. An alternative hypothesis of cellular transport of lysosomal enzymes in fibroblasts. Effect of inhibitors of lysosomal enzyme endocytosis on intra- and extra-cellular lysosomal enzyme activities 943–950
VON FIGURA, K. *see also* PRINZ, R. 671–676
VOS, J. *see* QUIRK, S. J. 159–167

WAHLE, K. W. J. *see* SCAIFE, J. R. 799–804

WALKER, J. H., BURGESS, R. J. & MAYER, R. J. Relative rates of turnover of subunits of mitochondrial proteins 927–932
WALKER, J. H. *see also* BURGESS, R. J. 919–926
WALKER, V. & TAYLOR, W. H. Ovalbumin digestion by human pepsins 1, 3 and 5 429–432
WALLIS, M. *see* BETTERIDGE, A. 319–323
WANG-IVERSON, P. *see* IVERSON, D. B. 175–178
WEBER, E. *see* VON FIGURA, K. 943–950
WHITE, D. A. *see* SPEAKE, B. K. 993–1000
WHITE, R. *see* SCHLESINGER, P. H. 103–109
WHITTON, P. D., RODRIGUES, L. M. & HEMS, D. A. Stimulation by vasopressin, angiotensin and oxytocin of gluconeogenesis in hepatocyte suspensions 893–898
WIGHTMAN, P. D. *see* BONNEY, R. J. 433–442
WILLIAMS, D. J. *see* PALMER, D. N. 9–14
WILLIAMS, J. F., BLACKMORE, P. F. & CLARK, M. G. New reaction sequences for the non-oxidative pentose phosphate pathway 257–282
WILLIAMS, J. F., CLARK, M. G. & BLACKMORE, P. F. The fate of ^{14}C in glucose 6-phosphate synthesized from [1-^{14}C]ribose 5-phosphate by enzymes of rat liver 241–256
WILLIAMS, T. *see* NEWSHOLME, E. A. 623–626
WILLIAMS-ASHMAN, H. G. *see* AHMED, K. 739–750
WILLIAMSON, D. H. *see* BENITO, M. 331–334; FERRÉ, P. 335–338, 759–765; ROBINSON, A. M. 343–346, 635–638
WILSON, M. J. *see* AHMED, K. 739–750
WILSON, S. B. Cyanide-insensitive oxidation of ascorbate + *NNN'N'*-tetramethyl-*p*-phenylenediamine mixture by mung-bean (*Phaseolus aureus*) mitochondria. An energy-linked function 129–136

YAMAMOTO, K. *see* FUNAKAWA, S. 977–981
YAMAMOTO, Y. *see* TOMIDA, M. 665–669

ZALESKI, J. & BRYŁA, J. Effect of alloxan-diabetes on gluconeogenesis and ureogenesis in isolated rabbit liver cells 563–568
ZAMBONI, M. *see* MONTANARO, L. 371–379
ZAPPIA, V. *see* DE ROSA, M. 1–7
ZRIKE, J. *see* TUAN, R. S. 67–74

Index of Subjects

2-Acetamidofluorene, evidence for the presence of a subpopulation of rat liver membrane-bound ribosomes that are detached by, and other carcinogens and by centrifugation (Palmer, D. N., Rabin, B. R. & Williams, D. J.) 9–14

Acetate, effect of, on the rate of biosynthesis of isocitrate lyase in *Chlorella fusca* var. *vacuolata* (Dunham, S. M. & Thurston, C. F.) 179–185

Acetoacetyl-coenzyme A deacylase, changes in the activities of 3-hydroxy-3-methylglutaryl-coenzyme A synthase and, in rat brain during postnatal development (Patel, T. B. & Clark, J. B.) 951–958

N-2-Acetylaminofluorene, *see* 2-Acetamidofluorene

N-Acetylglucosamine residues, terminal, role of rat liver non-parenchymal cells in the clearance from plasma of glycoproteins with, and mannose residues (Schlesinger, P. H., Doebber, T. W., Mandell, B. F., White, R., DeSchryver, C., Rodman, J. S., Miller, M. J. & Stahl, P.) 103–109

Adenine nucleotides, systematic variations in the concentrations of, and other purine nucleotides in steady-state perfused rat heart (Bates, D. J., Perrett, D. & Mowbray, J.) 485–493

Adenocarcinoma R3230AC cells, mammary-gland, rat, sodium ion-dependent and -independent transport of 2-aminoisobutyrate into, and the effects of streptozotocin-induced diabetes in the host animal (Hissin, P. J. & Hilf, R.) 205–215

Adenosine 3′:5′-cyclic monophosphate, 6-*N*,2′-*O*-dibutyryl-, effects of, and other agents on the biosynthesis and release of amylase by rat parotid gland (McPherson, M. A. & Hales, C. N.) 855–863

Adenosine 3′:5′-cyclic monophosphate, 6-*N*,2′-*O*-dibutyryl-, effects of glucagon and, on the transport of calcium ions by isolated rat liver mitochondria (Hughes, B. P. & Barritt, G. J.) 295–304

Adenosine 3′:5′-cyclic monophosphate, effects of angiotensin II and [1-sarcosine,8-alanine]angiotensin II on the production of, and the biosynthesis of steroids by isolated rat adrenal-gland-cortex glomerulosa cells (Bing, R. F. & Schulster, D.) 39–45

Adenosine 3′:5′-cyclic monophosphate, evidence for the unidirectional control by, of the function of human platelets (Haslam, R. J., Davidson, M. M. L. & Desjardins, J. V.) 83–95

Adenosine 3′:5′-cyclic monophosphate phosphodiesterase, effect of the redox state of the cytosolic nicotinamide–adenine dinucleotide couple on the activity of, and its role in the control of the responsiveness of isolated rat hepatocytes to glucagon (Clark, M. G. & Jarrett, I. G.) 805–816

Adenosine 3′:5′-cyclic monophosphate, use of obelin in a study of the inhibition by calcium ions of the formation of, in sealed pigeon erythrocyte 'ghosts' (Campbell, A. K. & Dormer, R. L.) 53–66

Adenosine, inhibition by analogues of, of the activity of adenylate cyclase in preparations of broken and intact human platelets (Haslam, R. J., Davidson, M. M. L. & Desjardins, J. V.) 83–95

Adenosine, opposite effects of starvation on the oxidation of, and on adenosine-induced release of insulin by isolated mouse islets of Langerhans (Andersson, A.) 619–621

Adenosine triphosphatase, coupling, mitochondrial, heart, ox, thermodynamic analysis of the interaction between, and its naturally occurring inhibitor protein (Gomez-Fernandez, J. C. & Harris, D. A.) 967–975

Adenylate cyclase, inhibition by adenosine analogues of the activity of, in preparations of broken and intact human platelets (Haslam, R. J., Davidson, M. M. L. & Desjardins, J. V.) 83–95

Adenylate cyclase, role of, in the effects of cholera toxin and guanosine [$\beta\gamma$-imido]triphosphate on the affinity of the β-adrenergic receptors of toad erythrocyte membranes (Fischer, J. & Sharp, G. W. G.) 505–510

Adipocytes, epididymal, rat, isolated, modulation by thyroid hormones of the number of β-adrenergic receptors in membranes of (Giudicelli, Y.) 1007–1010

Adipocytes, rat, isolated, stimulation by human very-low-density plasma lipoproteins of the biosynthesis of triacylglycerols in (Thomopoulos, P., Berthelier, M., Lagrange, D., Chapman, M. J. & Laudat, M.-H.) 169–174

Adipose tissue, epididymal, rat, inactivation of glycerol phosphate acyltransferase of, catalysed by adenosine 3′:5′-cyclic monophosphate-dependent protein kinase and its possible involvement in the regulation of the biosynthesis of triacylglycerols (Nimmo, H. G. & Houston, B.) 607–610

Adipose tissue, epididymal, rat, post-translational regulation of the activity of lipoprotein lipase in (Ashby, P., Bennett, D. P., Spencer, I. M. & Robinson, D. S.) 865–872

Adipose tissue, inguinal, sheep, utilization of methylmalonate for the biosynthesis of branched-chain fatty acids by preparations of (Scaife, J. R., Wahle, K. W. J. & Garton, G. A.) 799–804

Adrenal gland, rat, effect of the administration of corticotropin on the biosynthesis of a specific cytosolic protein in (Dazord, A., Gallet, D. & Saez, J. M.) 233–239

Adrenal-gland cortex, rat, effects of angiotensin II and [1-sarcosine,8-alanine]angiotensin II on the production of adenosine 3′:5′-cyclic monophosphate and the biosynthesis of steroids by glomerulosa cells isolated from (Bing, R. F. & Schulster, D.) 39–45

Adrenaline, effects of, and other agents on the biosynthesis and release of amylase by rat parotid gland (McPherson, M. A. & Hales, C. N.) 855–863

Adrenaline, regulation by, of the activity of phosphorylase phosphatase in rat skeletal muscle (Tao, S.-H., Huang, F. L., Lynch, A. & Glinsmann, W. H.) 347–350

Adrenaline, role of, in the post-translation regulation of the activity of lipoprotein lipase in rat epididymal adipose tissue (Ashby, P., Bennett, D. P., Spencer, I. M. & Robinson, D. S.) 865–872

α-Adrenergic agonists, dependence on calcium ions of the inhibition by cholinergic agonists and, of the biosynthesis of protein in rat parotid-gland slices (Kanagasuntheram, P. & Lim, S. C.) 23–29

β-Adrenergic receptors, membrane, erythrocyte, toad, effects of cholera toxin and guanosine [βγ-imido]triphosphate on the affinity of (Fischer, J. & Sharp, G. W. G.) 505–510

β-Adrenergic receptors, modulation by thyroid hormones of the number of, in membranes of isolated rat epididymal adipocytes (Guidicelli, Y.) 1007–1010

Adrenocorticotropin, *see* Corticotropin

Age, changes in the composition and structure of human articular-cartilage proteoglycans with (Bayliss, M. T. & Ali, S. Y.) 683–693

Age, variation of the concentrations of carnitine in various rat tissues with sex and (Borum, P. R.) 677–681

Alanine, accelerated degradation of protein formed in rat and rabbit reticulocytes under conditions of depletion of, and other individual amino acids (Chandler, C. S. & Ballard, F. J.) 151–158

Alanine, characteristics of the transport of serine, glutamine and, across the plasma membrane of isolated rat hepatocytes (Joseph, S. K., Bradford, N. M. & McGivan, J. D.) 827–836

Aldosterone, effects of angiotensin II and [1-sarcosine,8-alanine]angiotensin II on the production of adenosine 3′:5′-cyclic monophosphate and the biosynthesis of, and other steroids by isolated rat adrenal-gland-cortex glomerulosa cells (Bing, R. F. & Schulster, D.) 39–45

Alloxan-diabetes, *see* Diabetes, alloxan-induced

Amino acid, labelled, analysis of errors in estimation of the rate of biosynthesis of protein by constant infusion of, into an animal (Garlick, P. J.) 402–405

Amino acids, characteristics of the transport of, across the plasma membrane of isolated rat hepatocytes (Joseph, S. K., Bradford, N. M. & McGivan, J. D.) 827–836

Amino acids, individual, accelerated degradation of protein formed in rat and rabbit reticulocytes under conditions of depletion of (Chandler, C. S. & Ballard, F. J.) 151–158

Amino acids, role of phosphoenolpyruvate carboxykinase in the metabolism of, in skeletal muscle and heart muscle of various animal species (Newsholme, E. A. & Williams, T.) 623–626

2-Aminoisobutyrate, sodium ion-dependent and -independent transport of, into rat mammary-gland adenocarcinoma R3230AC cells and the effects of streptozotocin-induced diabetes in the host animal (Hissin, P. J. & Hilf, R.) 205–215

3-Aminopicolinate, induction of hyperglycaemia and other metabolic effects of administration of, to the rat (MacDonald, M. J., Huang, M.-T. & Lardy, H. A.) 495–504

Ammonia, sources of, for the biosynthesis of urea by isolated rat hepatocytes (Krebs, H. A., Hems, R., Lund, P., Halliday, D. & Read, W. W. C.) 733–737

Ammonium chloride, effects of bicarbonate and, on the activity of glutaminase in isolated rat liver mitochondria (Joseph, S. K. & McGivan, J. D.) 837–844

Amylase, control of the biosynthesis and release of, by rat parotid gland (McPherson, M. A. & Hales, C. N.) 855–863

α-Amylase, sensitivity towards proteinases of the release of ribonuclease and, by rat pancreatic microsomal fraction (Pearce, P. D., May, B. K. & Elliott, W. H.) 611–614

Angiotensin, stimulation by vasopressin, oxytocin and, of gluconeogenesis in isolated rat hepatocytes (Whitton, P. D., Rodrigues, L. M. & Hems, D. A.) 893–898

Angiotensin II, effects of [1-sarcosine,8-alanine]angiotensin II and, on the production of adenosine 3′:5′-cyclic monophosphate and the biosynthesis of steroids by isolated rat adrenal-gland-cortex glomerulosa cells (Bing, R. F. & Schulster, D.) 39–45

Angiotensin II, [1-sarcosine,8-alanine]-, effects of angiotensin II and, on the production of adenosine 3′:5′-cyclic monophosphate and the biosynthesis of steroids by isolated rat adrenal-gland-cortex glomerulosa cells (Bing, R. F. & Schulster, D.) 39–45

Anions, ratios of, and calcium ions and the production of protons during the uptake of calcium ions by rat heart and liver mitochondria (Harris, E. J.) 983–991

Artemia salina, *see* Brine shrimp

Ascites cells, Krebs II, characterization of the acidic ribosomal phosphoprotein L_7 from, and its relationship to *Escherichia coli* ribosomal proteins L7 and L12 (Leader, D. P. & Coia, A. A.) 569–572

Ascorbate, cyanide-insensitive uptake of oxygen by mung-bean mitochondria in the presence of a mixture of, and *NNN′N′*-tetramethyl-*p*-phenylenediamine as electron donor (Wilson, S. B.) 129–136

Aspartate aminotransferase, evidence for the participation of, in the formation of glucose in liver of suckling newborn rats (Ferré, P. & Williamson, D. H.) 335–338

Aspergillus ochraceus TS, specific 11α-hydroxylation of progesterone by (Samanta, T. B., Roy, N. & Chattopadhyay, S.) 593–594

Bacillus megaterium KM, accumulation of calcium ions during sporulation of (Hogarth, C. & Ellar, D. J.) 197–203

Bacillus subtilis, inhibition by pentachlorophenol and related uncouplers of oxidative phosphorylation of the transport of proline and glycine by (Nicholas, R. A. & Ordal, G. W.) 639–647

Bean, broad (*Vicia faba*), formation of isomers of biliverdin during the degradation of leghaemoglobins from root nodules of, and other leguminous plants (Lehtovaara, P. & Perttilä, U.) 359–364

Bean, kidney (*Phaseolus vulgaris*), formation of isomers of biliverdin during the degradation of leghaemoglobins from root nodules of, and other leguminous plants (Lehtovaara, P. & Perttilä, U.) 359–364

Bean, kidney (*Phaseolus vulgaris*), oxidation of glycine by leghaemoglobin from root nodules of, and associated catabolic reactions at the haem group (Lehtovaara, P.) 351–358

Bean, mung (*Phaseolus aureus*), cyanide-insensitive uptake of oxygen by mitochondria from, in the presence of ascorbate/*NNN′N′*-tetramethyl-*p*-phenylenediamine mixture as electron donor (Wilson, S. B.) 129–136

Bean, soya (*Glycine max*), formation of isomers of biliverdin during the degradation of leghaemoglobins from root nodules of, and other leguminous plants (Lehtovaara, P. & Perttilä, U.) 359–364

BHK cells, *see* Cells, kidney, baby-hamster

Bicarbonate, effects of ammonium chloride and, on the activity of glutaminase in isolated rat liver mitochondria (Joseph, S. K. & McGivan, J. D.) 837–844

Bile, excretion of cyclohexylphenyl 4-sulphate in, in the guinea pig (Powell, G. M., Olavesen, A. H. & Curtis, C. G.) 443–448
Bile pigments, formation of, during the degradation of root-nodule leghaemoglobins from various leguminous plants (Lehtovaara, P. & Perttilä, U.) 359–364
Biliverdin, formation of isomers of, during the degradation of root-nodule leghaemoglobins from various leguminous plants (Lehtovaara, P. & Perttilä, U.) 359–364
Blood platelets, human, broken and intact, inhibition by adenosine analogues of the activity of adenylate cyclase in preparations of (Haslam, R. J., Davidson, M. M. L. & Desjardins, J. V.) 83–95
Blood platelets, human, evidence that the formation of malondialdehyde in, is catalysed by thromboxane synthase (McMillan, R. M., MacIntyre, D. E., Booth, A. & Gordon, J. L.) 595–598
Blood, rat, changes in the concentrations of ketone bodies in, during postnatal development (Ferré, P., Pégorier, J.-P., Williamson, D. H. & Girard, J. R.) 759–765
Blood, rat, increased concentration of glucose in, and other metabolic effects of administration of 3-aminopicolinate to the animals (MacDonald, M. J., Huang, M.-T. & Lardy, H. A.) 495–504
Blood, rat, role of the liver non-parenchymal cells in the clearance from, of glycoproteins with terminal mannose and *N*-acetylglucosamine residues (Schlesinger, P. H., Doebber, T. W., Mandell, B. F., White, R., DeSchryver, C., Rodman, J. S., Miller, M. J. & Stahl, P.) 103–109
Brain, guinea-pig, activation by calcium ions of the hydrolysis of phosphatidyl-*myo*-inositol 4-phosphate (diphosphoinositide) and phosphatidyl-*myo*-inositol 4,5-bisphosphate (triphosphoinositide) in synaptosomes from (Griffin, H. D. & Hawthorne, J. N.) 541–552
Brain, rat, changes in the activities of acetoacetyl-coenzyme A deacylase and 3-hydroxy-3-methylglutaryl-coenzyme A synthase in, during postnatal development (Patel, T. B. & Clark, J. B.) 951–958
Brain, rat, female, metabolism of high-molecular-weight ribonucleic acid in hypothalamic and cortical regions of, during development (Hall, C. & Lim, L.) 511–521
Brain, rat, hydrolysis of phosphatidylinositol by lysosomal enzymes of, and liver (Irvine, R. F., Hemington, N. & Dawson, R. M. C.) 475–484
Brain, rat, rapid method for the isolation of a relatively pure metabolically competent preparation of synaptosomes from (Booth, R. F. G. & Clark, J. B.) 365–370
Brine shrimp (*Artemia salina*), inhibitory effect of modeccin on the elongation of polypeptide chains during the biosynthesis of protein catalysed by ribosomes from (Montanaro, L., Sperti, S., Zamboni, M., Denaro, M., Testoni, G., Gasperi-Campani, A. & Stirpe, F.) 371–379
Broad bean, *see* Bean, broad
Bufo marinus, *see* Toad

Calcitonin, possible role of 5-hydroxytryptamine as mediator in the action of, on rat gastrointestinal tract and pancreas (Nakhla, A. M. & Latif, A.) 339–342
Calcium ions, accumulation of, during sporulation of *Bacillus megaterium* KM (Hogarth, C. & Ellar, D. J.) 197–203
Calcium ions, activation by, of the hydrolysis of phosphatidyl-*myo*-inositol 4-phosphate (diphosphoinositide) and phosphatidyl-*myo*-inositol 4,5-bisphosphate (triphosphoinositide) by guinea-pig brain synaptosomes (Griffin, H. D. & Hawthorne, J. N.) 541–552
Calcium ions, anion/calcium ion ratios and production of protons during the uptake of, by rat heart and liver mitochondria (Harris, E. J.) 983–991
Calcium ions, characterization of human erythrocyte membrane proteins after cell fusion induced by oleoylglycerol in the presence of (Quirk, S. J., Ahkong, Q. F., Botham, G. M., Vos, J. & Lucy, J. A.) 159–167
Calcium ions, dependence on, of the inhibition by α-adrenergic agonists and cholinergic agonists of the biosynthesis of protein in rat parotid-gland slices (Kanagasuntheram, P. & Lim, S. C.) 23–29
Calcium ions, effects of glucagon and 6-*N*,2′-*O*-dibutyryladenosine 3′:5′-cyclic monophosphate on the transport of, by isolated rat liver mitochondria (Hughes, B. P. & Barritt, G. J.) 295–304
Calcium ions, functional involvement of carbonic anhydrase in the transport of, by chick-embryo chorioallantoic membrane (Tuan, R. S. & Zrike, J.) 67–74
Calcium ions, regulation of the extramitochondrial concentration of, by rat liver mitochondria (Nicholls, D. G.) 463–474
Calcium ions, role of, in the regulation of the activity of nicotinamide–adenine dinucleotide-linked isocitrate dehydrogenase in mitochondria from rat heart and other tissues (Denton, R. M., Richards, D. A. & Chin, J. G.) 899–906
Calcium ions, stable enhancement of the retention of, by isolated rat liver mitochondria after the administration of glucagon to the animal (Prpić, V., Spencer, T. L. & Bygrave, F. L.) 705–714
Calcium ions, use of obelin in a study of the inhibition by, of the formation of adenosine 3′:5′-cyclic monophosphate in sealed pigeon erythrocyte 'ghosts' (Campbell, A. K. & Dormer, R. L.) 53–66
Caldariella acidophila, pathway for the biosynthesis of polyamines in (De Rosa, M., De Rosa, S., Gambacorta, A., Carteni-Farina. M. & Zappia, V.) 1–7
Carbamoylcholine, effects of, and other agents on the biosynthesis and release of amylase by rat parotid gland (McPherson, M. A. & Hales, C. N.) 855–863
Carbon isotope (^{14}C), distribution of, in glucose 6-phosphate biosynthesized from [1-^{14}C]ribose 5-phosphate by a rat liver extract and its relevance to the pentose phosphate pathway for the metabolism of glucose (Williams, J. F., Clark, M. G. & Blackmore, P. F.) 241–256
Carbonic anhydrase, functional involvement of, in the transport of calcium ions by chick-embryo chorioallantoic membrane (Tuan, R. S. & Zrike, J.) 67–74
Carnitine, variation with age and sex of the concentrations of, in various rat tissues (Borum, P. R.) 677–681
Carotenoids, protective role of, in the photosensitivity of respiration by *Neurospora crassa* mitochondria (Ramadan-Talib, Z. & Prebble, J.) 767–775
Cartilage, articular, human, age-related changes in the composition and structure of proteoglycans from (Bayliss, M. T. & Ali, S. Y.) 683–693

Cells, adenocarcinoma R3230AC, mammary-gland, rat, sodium ion-dependent and -independent transport of 2-aminoisobutyrate into, and the effects of streptozotocin-induced diabetes in the host animal (Hissin, P. J. & Hilf, R.) 205–215

Cells, ascites, Krebs II, characterization of the acidic ribosomal phosphoprotein L_y from, and its relationship to *Escherichia coli* ribosomal proteins L7 and L12 (Leader, D. P. & Coia, A. A.) 569–572

Cells, hepatoma, rat, cultured, effect of glutamine on the degradation of glutamine synthetase in, derived from Morris hepatoma 7288C (Crook, R. B. & Tomkins, G. M.) 47–52

Cells, hepatoma, rat, cultured, extensive conjugation of metabolites of 3,4-dihydroxyphenethylamine in, and human skin fibroblasts (Crooks, P. A., Breakefield, X. O., Sulens, C. H., Castiglione, C. M. & Coward, J. K.) 187–196

Cells, kidney (BHK-21/C13), baby-hamster, onset of the degradation of ribosomes during cessation of growth in (Melvin, W. T. & Keir, H. M.) 933–941

Cells, Leydig, testis, mouse, induction by the β-subunit of human chorionic gonadotropin of the production of testosterone by (Ramakrishnan, S., Das, C. & Talwar, G. P.) 599–602

Cells, neuroblastoma, mouse, cultured, regulation of the biosynthesis of guanosine 3′:5′-cyclic monophosphate in (Bartfai, I., Breakefield, X. O. & Greengard, P.) 119–127

Cells, neuroblastoma, mouse, method for the measurement of relative changes in the concentration of guanosine 3′:5′-cyclic monophosphate in, on muscarinic cholinergic stimulation (Strange, P. G.) 583–590

Cells, neuroblastoma, N2a, mouse, growth and the metabolism of plasma-membrane fucosylated glycoproteins in (Milenkovic, A. G., Rachmeler, M. & Johnson, T. C.) 695–704

Cells, *see also* Fibroblasts

Cerebral cortex, rat, female, metabolism of high-molecular-weight ribonucleic acid in, and hypothalamus during development (Hall, C. & Lim, L.) 511–521

Chick embryo, effects of oestrogens on the activity of ornithine decarboxylase and the concentrations of polyamines in differentiating Müllerian ducts and other organs of (Teng, C. S. & Teng, C. T.) 143–149

Chick embryo, functional involvement of carbonic anhydrase in the transport of calcium ions by chorioallantoic membrane from (Tuan, R. S. & Zrike, J.) 67–74

Chick, metabolism of 1α-hydroxycholecalciferol in (Edelstein, S., Noff, D., Freeman, D., Sheves, M. & Mazur, Y.) 111–117

Chick, turnover of protein in skeletal muscle of, during growth (Maruyama, K., Sunde, M. L. & Swick, R. W.) 573–582

Chicken, changes in the rates of biosynthesis and degradation of protein in skeletal muscle of, during stretch-induced hypertrophy of skeletal muscle (Laurent, G. J., Sparrow, M. P. & Millward, D. J.) 407–417

Chicken, content and turnover of collagen in skeletal muscle and heart muscle of, during stretch-induced hypertrophy of skeletal muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 419–427

Chicken, isolation of a high-molecular-weight high-mobility-group-type non-histone protein from oviduct from (Teng, C. S., Gallagher, K. & Teng, C. T.) 1003–1006

Chicken, rates of biosynthesis of protein in skeletal muscle, heart muscle and smooth muscle of (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 393–401

Chicken, utilization of methylmalonate for the biosynthesis of branched-chain fatty acids by preparations of liver from (Scaife, J. R., Wahle, K. W. J. & Garton, G. A.) 799–804

Chlorate, characterization of a mutant of *Escherichia coli* K12 that is sensitive to, when grown aerobically (Giordano, G., Grillet, L., Rosset, R., Dou, J. H., Azoulay, E. & Haddock, B. A.) 553–561

Chlorella fusca var. *vacuolata*, effect of acetate on the rate of biosynthesis of isocitrate lyase in (Dunham, S. M. & Thurston, C. F.) 179–185

[7-^{3}H]Cholecalciferol, 1α-hydroxy-, synthesis of, and its metabolism in the chick (Edelstein, S., Noff, D., Freeman, D., Sheves, M. & Mazur, Y.) 111–117

Cholera toxin, effects of guanosine [βγ-imido]triphosphate and, on the affinity of β-adrenergic receptors of toad erythrocyte membranes (Fischer, J. & Sharp, G. W. G.) 505–510

Choline kinase, evidence from essential-fatty acid-deficient animals for a regulatory role of, in the biosynthesis of phosphatidylcholine in rat liver (Infante, J. P. & Kinsella, J. E.) 631–633

Cholinergic agonists, dependence on calcium ions of the inhibition by α-adrenergic agonists and, of the biosynthesis of protein in rat parotid-gland slices (Kanagasuntheram, P. & Lim, S. C.) 23–29

Cholinergic stimulation, muscarinic, method for the measurement of relative changes in the concentration of guanosine 3′:5′-cyclic monophosphate in mouse neuroblastoma cells on (Strange, P. G.) 583–590

Chorioallantoic membrane, chick-embryo, functional involvement of carbonic anhydrase in the transport of calcium ions by (Tuan, R. S. & Zrike, J.) 67–74

Chorionic gonadotropin, *see* Gonadotropin, chorionic

Chromatin, effects of polyamines on the reactions catalysed by protein kinases associated with non-histone proteins and, of rat ventral-prostate-gland nuclei (Ahmed, K., Wilson, M. J., Goueli, S. A. & Williams-Ashman, H. G.) 739–750

Chromatin, hypothalamus, sheep, binding of oestradiol-17β-receptor protein complex and of progesterone-receptor protein complex to, from male and female animals (Perry, B. N. & Lopez, A.) 873–883

Clearing-factor lipase, *see* Lipoprotein lipase

Coenzyme B_{12} (5′-deoxyadenosylcobalamin), concerted induction by ethanolamine and, of the formation of ethanolamine ammonia-lyase in *Escherichia coli* (Blackwell, C. M. & Turner, J. M.) 751–757

Collagen, biosynthesis and partial characterization of protein components similar to, of rat lens capsule (Heathcote, J. G., Sear, C. H. J. & Grant, M. E.) 283–294

Collagen, content and turnover of, in chicken skeletal muscle and heart muscle during stretch-induced hypertrophy of skeletal muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 419–427

Coproporphyrinogen III oxidase, location of the activity of, in rat liver mitochondria (Grandchamp, B., Phung, N. & Nordmann, Y.) 97–102
Corticosterone, effects of angiotensin II and [1-sarcosine,-8-alanine]angiotensin II on the production of adenosine 3′:5′-cyclic monophosphate and the biosynthesis of, and other steroids by isolated rat adrenal-gland-cortex glomerulosa cells (Bing, R. F. & Schulster, D.) 39–45
Corticotropin, effect of the administration of, on the biosynthesis of a specific cytosolic protein in rat adrenal gland (Dazord, A., Gallet, D. & Saez, J. M.) 233–239
Corticotropin, effects of cortisol, thyroxine and, on the biosynthesis of glycerolipids and on the activity of phosphatidate phosphatase in rat liver (Glenny, H. P. & Brindley, D. N.) 777–784
Cortisol, effects of corticotropin, thyroxine and, on the biosynthesis of glycerolipids and on the activity of phosphatidate phosphatase in rat liver (Glenny, H. P. & Brindley, D. N.) 777–784
Cyclic adenosine 3′:5′-monophosphate, *see* Adenosine 3′:5′-cyclic monophosphate
Cyclic guanosine 3′:5′-monophosphate, *see* Guanosine 3′:5′-cyclic monophosphate
Cyclic nucleotide phosphodiesterases, properties of the calcium ion-dependent protein modulator of the activities of, of mouse epidermis (Murray, A. W. & Rogers, A.) 727–732
Cyclohexylphenyl 4-sulphate, biliary excretion of, in the guinea pig (Powell, G. M., Olavesen, A. H. & Curtis, C. G.) 443–448
Cysteine, participation of cystine and, in the inactivation of tyrosine aminotransferase in rat liver homogenates (Buckley, W. T. & Milligan, L. P.) 449–454
Cystine, participation of cysteine and, in the inactivation of tyrosine aminotransferase in rat liver homogenates (Buckley, W. T. & Milligan, L. P.) 449–454
Cytoplasm, liver, rat, evidence that most of the membrane proteins of rat liver lysosomes are exposed to (Schneider, D. L., Burnside, J., Gorga, F. R. & Nettleton, C. J.) 75–82
Cytosol, adrenal-gland, rat, effect of the administration of corticotropin on the biosynthesis of a specific protein in (Dazord, A., Gallet, D. & Saez, J. M.) 233–239
Cytosol, uterus, human, interaction of progestins with the progesterone-receptor protein of (Kasid, A., Buckshee, K., Hingorani, V. & Laumas, K. R.) 531–539
Cytosol, uterus, rat, binding characteristics, content and distribution of oestrogen-receptor proteins in, and nuclei in relation to intra-uterine devices and the oestrous cycle (Myatt, L., Chaudhuri, G., Elder, M. G. & Lim, L.) 523–529

5′-Deoxyadenosylcobalamin (coenzyme B_{12}), concerted induction by ethanolamine and, of the formation of ethanolamine ammonia-lyase in *Escherichia coli* (Blackwell, C. M. & Turner, J. M.) 751–757
Deoxyribonucleic acid, initiation of the transcription of, for the biosynthesis of ribonucleic acid in normal and phosphorylated rat liver nucleosomes (Ord, M. G. & Stocken, L. A.) 615–618
Development, adult, changes in the composition and structure of human articular-cartilage proteoglycans during (Bayliss, M. T. & Ali, S. Y.) 683–693
Development, metabolism of high-molecular-weight ribonucleic acid in hypothalamic and cortical regions of female rat brain during (Hall, C. & Lim, L.) 511–521
Development, postnatal, changes in the activities of acetoacetyl-coenzyme A deacylase and 3-hydroxy-3-methylglutaryl-coenzyme A synthase during (Patel, T. B. & Clark, J. B.) 951–958
Development, postnatal, regulation of ketogenesis in the rat during (Ferré, P., Pégorier, J.-P., Williamson, D. H. & Girard, J. R.) 759–765
Development, postnatal, turnover of protein in chick skeletal muscle during (Maruyama, K., Sunde, M. L. & Swick, R. W.) 573–582
Development, postnatal, variation of the concentrations of carnitine in various rat tissues with sex and during (Borum, P. R.) 677–681
Diabetes, alloxan-induced, effect of, on gluconeogenesis and ureogenesis in isolated rabbit liver cells (Zaleski, J. & Bryła, J.) 563–568
Diabetes, streptozotocin-induced, effects of, in the host animal on the sodium ion-dependent and -independent transport of 2-aminoisobutyrate into rat mammary-gland adenocarcinoma R3230AC cells (Hissin, P. J. & Hilf, R.) 205–215
Diamide, role of reduced glutathione in the reduction of, by isolated rat liver mitochondria (Jocelyn, P. C.) 649–664
11,11-Diamino-4,8-diazaundecane (sym-nor-spermine), pathway for the biosynthesis of, and other polyamines in *Caldariella acidophila* (De Rosa, M., De Rosa, S., Gambacorta, A., Cartenì-Farina, M. & Zappia, V.) 1–7
6-*N*,2′-*O*-Dibutyryladenosine 3′:5′-cyclic monophosphate, effects of, and other agents on the biosynthesis and release of amylase by rat parotid gland (McPherson, M. A. & Hales, C. N.) 855–863
6-*N*,2′-*O*-Dibutyryladenosine 3′:5′-cyclic monophosphate, effects of glucagon and, on the transport of calcium ions by isolated rat liver mitochondria (Hughes, B. P. & Barritt, G. J.) 295–304
3,4-Dihydroxyphenethylamine (dopamine), extensive conjugation of metabolites of, in cultured human skin fibroblasts and rat hepatoma cells (Crooks, P. A., Breakefield, X. O., Sulens, C. H., Castiglione, C. M. & Coward, J. K.) 187–196
Diphosphoinositide, *see* Phosphatidyl-*myo*-inositol 4-phosphate
Dipicolinate, role of, in the accumulation of calcium ions during sporulation of *Bacillus megaterium* KM (Hogarth, C. & Ellar, D. J.) 197–203
Dopamine (3,4-dihydroxyphenethylamine), extensive conjugation of metabolites of, in cultured human skin fibroblasts and rat hepatoma cells (Crooks, P. A., Breakefield, X. O., Sulens, C. H., Castiglione, C. M. & Coward, J. K.) 187–196

Electrons, cyanide-insensitive uptake of oxygen by mung-bean mitochondria in the presence of ascorbate/*NNN′N′*-tetramethyl-*p*-phenylenediamine mixture as donor of (Wilson, S. B.) 129–136

Endocytosis, effect of inhibitors of, on the intracellular and extracellular activities of lysosomal enzymes of fibroblasts (von Figura, K. & Weber, E.) 943–950
Endoplasmic reticulum, *see* Reticulum, endoplasmic
Enzymes, lysosomal, effect of inhibitors of endocytosis on the intracellular and extracellular activities of, of fibroblasts (von Figura, K. & Weber, E.) 943–950
Epidermis, mouse, properties of the calcium ion-dependent protein modulator of the activities of cyclic nucleotide phosphodiesterases of (Murray, A. W. & Rogers, A.) 727–732
Epididymis, rat, inactivation of glycerol phosphate acyltransferase of adipose tissue of, catalysed by adenosine 3′:5′-cyclic monophosphate-dependent protein kinase and its possible involvement in the regulation of the biosynthesis of triacylglycerols (Nimmo, H. G. & Houston, B.) 607–610
Epididymis, rat, modulation by thyroid hormones of the number of β-adrenergic receptors in membranes of adipocytes isolated from (Giudicelli, Y.) 1007–1010
Epididymis, rat, post-translational regulation of the activity of lipoprotein lipase in adipose tissue from (Ashby, P., Bennett, D. P., Spencer, I. M. & Robinson, D. S.) 865–872
Epinephrine, *see* Adrenaline
Erythrocyte 'ghosts', pigeon, sealed, use of obelin in a study of the inhibition by calcium ions of the formation of adenosine 3′:5′-cyclic monophosphate in (Campbell, A. K. & Dormer, R. L.) 53–66
Erythrocytes, human, characterization of membrane proteins of, after cell fusion induced by oleoylglycerol in the presence of calcium ions (Quirk, S. J., Ahkong, Q. F., Botham, G. M., Vos, J. & Lucy, J. A.) 159–167
Erythrocytes, toad, effects of cholera toxin and guanosine 5′-[βγ-imido]triphosphate on the affinity of the β-adrenergic receptors of membranes of (Fischer, J. & Sharp, G. W. G.) 505–510
Escherichia coli, characterization of the acidic ribosomal phosphoprotein L_7 from Krebs II ascites cells and its relationship to ribosomal proteins L7 and L12 from (Leader, D. P. & Coia, A. A.) 569–572
Escherichia coli, concerted induction by ethanolamine and coenzyme B_{12} of the formation of ethanolamine ammonia-lyase in (Blackwell, C. M. & Turner, J. M.) 751–757
Escherichia coli K12, characterization of a mutant of, that is sensitive to chlorate when grown aerobically (Giordano, G., Grillet, L., Rosset, R., Dou, J. H., Azoulay, E. & Haddock, B. A.) 553–561
Escherichia coli M.R.E. 600, inhibition by pseudomonic acid of the activity of isoleucyl-transfer ribonucleic acid synthetase in (Hughes, J. & Mellows, G.) 305–318
Estradiol, *see* Oestradiol
Estrogens, *see* Oestrogens
Estrous cycle, *see* Oestrous cycle
Ethanol, mechanisms of the effects of the administration of glycerol and, on the activity of phosphatidate phosphatase in rat liver (Savolainen, M. J. & Hassinen, I .E.) 885–892
Ethanolamine ammonia-lyase, concerted induction by ethanolamine and coenzyme B_{12} of the formation of, in *Escherichia coli* (Blackwell, C. M. & Turner, J. M.) 751–757
Ethanolamine, concerted induction by coenzyme B_{12} and, of the formation of ethanolamine ammonia-lyase in *Escherichia coli* (Blackwell, C. M. & Turner, J. M.) 751–757
Eye, rat, biosynthesis and partial characterization of collagenous protein components of the lens capsule of (Heathcote, J. G., Sear, C. H. J. & Grant, M. E.) 283–294

Fat-cells, epididymal, rat, isolated, modulation by thyroid hormones of the numbers of β-adrenergic receptors in membranes of (Giudicelli, Y.) 1007–1010
Fat-cells, rat, isolated, stimulation by human very-low-density plasma lipoproteins of the biosynthesis of triacylglycerols in (Thomopoulos, P., Berthelier, M., Lagrange, D., Chapman, M. J. & Laudat, M.-H.) 169–174
Fatty acids, branched-chain, utilization of methylmalonate for the biosynthesis of, by preparations of chicken liver and sheep inguinal adipose tissue (Scaife, J. R., Wahle, K. W. J. & Garton, G. A.) 799–804
Fatty acids, essential, effect of dietary deficiency of, on the activity of choline kinase in rat liver (Infante, J. P. & Kinsella, J. E.) 631–633
Fibroblasts, effect of inhibitors of endocytosis on the intracellular and extracellular activities of lysosomal enzymes of (von Figura, K. & Weber, E.) 943–950
Fibroblasts, skin, human, cultured, endocytosis of sulphated proteoglycans by (Prinz, R., Schwermann, J., Buddecke, E. & von Figura, K.) 671–676
Fibroblasts, skin, human, cultured, extensive conjugation of metabolites of 3,4-dihydroxyphenethylamine in, and rat hepatoma cells (Crooks, P. A., Breakefield, X. O., Sulens, C. H., Castiglione, C. M. & Coward, J. K.) 187–196
Frog (*Rana temporaria*), chemical-modification approach to elucidation of the olfactory code exemplified by the inhibition by mersalyl of the response of the olfactory mucosa of, to pentyl acetate and related odorants (Menevse, A., Dodd, G. & Poynder, T. M.) 845–854

Gastrointestinal tract, rat, possible role of 5-hydroxytryptamine as mediator in the action of calcitonin on, and pancreas (Nakhla, A. M. & Latif, A.) 339–342
Glomerulosa cells, adrenal-gland-cortex, rat, isolated, effects of angiotensin II and [1-sarcosine,8-alanine]-angiotensin II on the production of adenosine 3′:5′-cyclic monophosphate and the biosynthesis of steroids by (Bing, R. F. & Schulster, D.) 39–45
Glucagon, effect of the redox state of the cytosolic nicotinamide–adenine dinucleotide couple on the activity of adenosine 3′:5′-cyclic monophosphate and its role in the control of the responsiveness of isolated rat hepatocytes to (Clark, M. G. & Jarrett, I. G.) 805–816
Glucagon, effects of 6-*N*,2′-*O*-dibutyryladenosine 3′:5′-cyclic monophosphate and, on the transport of calcium ions by isolated rat liver mitochondria (Hughes, B. P. & Barritt, G. J.) 295–304
Glucagon, effects of phenylephrine, vasopressin and, on gluconeogenesis and on enzymes involved in the metabolism of glycogen in isolated rat hepatocytes (Hue, L., Felíu, J. E. & Hers, H.-G.) 791–797

Glucagon, stable enhancement of the retention of calcium ions by isolated rat liver mitochondria after the administration of, to the animal (Prpić, V., Spencer, T. L. & Bygrave, F. L.) 705–714

Gluconeogenesis, differential effects of tryptophan on, in isolated rat and guinea-pig hepatocytes (Smith, S. A., Elliott, K. R. F. & Pogson, C. I.) 817–825

Gluconeogenesis, effects of alloxan-induced diabetes on ureogenesis and, in isolated rabbit liver cells (Zaleski, J. & Bryła, J.) 563–568

Gluconeogenesis, effects of phenylephrine, vasopressin and glucagon on, and on enzymes involved in the metabolism of glycogen in isolated rat hepatocytes (Hue, L., Felíu, J. E. & Hers, H.-G.) 791–797

Gluconeogenesis, evidence for the participation of aspartate aminotransferase in, in liver of suckling newborn rats (Ferré, P. & Williamson, D. H.) 335–338

Gluconeogenesis, stimulation by vasopressin, angiotensin and oxytocin of, in isolated rat hepatocytes (Whitton, P. D., Rodrigues, L. M. & Hems, D. A.) 893–898

Glucose, differential effects of tryptophan on the formation of, in isolated rat and guinea-pig hepatocytes (Smith, S. A., Elliott, K. R. F. & Pogson, C. I.) 817–825

Glucose, distribution of ^{14}C in glucose 6-phosphate biosynthesized from [1-^{14}C]ribose 5-phosphate by a rat liver extract and its relevance to the pentose phosphate pathway for the metabolism of (Williams, J. F., Clark, M. G. & Blackmore, P. F.) 241–256

Glucose, effect of exogenous pyruvate on the induction by, of the release of insulin by isolated rat islets of Langerhans (Sener, A., Kawazu, S., Hutton, J. C., Boschero, A. C., Devis, G., Somers, G., Herchuelz, A. & Malaisse, W. J.) 217–232

Glucose, effects of alloxan-induced diabetes on the formation of urea and, in isolated rabbit liver cells (Zaleski, J. & Bryła, J.) 563–568

Glucose, effects of phenylephrine, vasopressin and glucagon on the formation of, and on enzymes involved in the metabolism of glycogen in isolated rat hepatocytes (Hue, L., Felíu, J. E. & Hers, H.-G.) 791–797

Glucose, evidence for new reaction sequences in the pentose phosphate pathway for the metabolism of, by a rat liver extract (Williams, J. F., Blackmore, P. F. & Clark, M. G.) 257–282

Glucose, evidence for the participation of aspartate aminotransferase in the formation of, in liver of suckling newborn rats (Ferré, P. & Williamson, D. H.) 335–338

Glucose, formation and characterization of lipid-linked oligosaccharides containing, of lactating rabbit mammary gland (Speake, B. K. & White, D. A.) 993–1000

Glucose, increased concentration of, in rat blood and other metabolic effects of administration of 3-aminopicolinate to the animals (MacDonald, M. J., Huang, M.-T. & Lardy, H. A.) 495–504

Glucose 6-phosphate, distribution of ^{14}C in, biosynthesized from [1-^{14}C]ribose 5-phosphate by a rat liver extract and its relevance to the pentose phosphate pathway for the metabolism of glucose (Williams, J. F., Clark, M. G. & Blackmore, P. F.) 241–256

Glucose, stimulation by vasopressin, angiotensin and oxytocin of the formation of, in isolated rat hepatocytes (Whitton, P. D., Rodrigues, L. M. & Hems, D. A.) 893–898

Glutaminase, effects of ammonium chloride and bicarbonate on the activity of, in isolated rat liver mitochondria (Joseph, S. K. & McGivan, J. D.) 837–844

Glutamine, characteristics of the transport of alanine, serine and, across the plasma membrane of isolated rat hepatocytes (Joseph, S. K., Bradford, N. M. & McGivan, J. D.) 827–836

Glutamine, effect of, on the degradation of glutamine synthetase in cultured rat hepatoma cells derived from Morris hepatoma 7288C (Crook, R. B. & Tomkins, G. M.) 47–52

Glutamine synthetase, effect of glutamine on the degradation of, in cultured rat hepatoma cells derived from Morris hepatoma 7288C (Crook, R. B. & Tomkins, G. M.) 47–52

Glutathione, reduced, role of, in the reduction of diamide by isolated rat liver mitochondria (Jocelyn, P. C.) 649–664

Glycerol, mechanisms of the effects of the administration of ethanol and, on the activity of phosphatidate phosphatase in rat liver (Savolainen, M. J. & Hassinen, I. E.) 885–892

Glycerol mono-oleate, characterization of human erythrocyte membrane proteins after cell fusion induced by, in the presence of calcium ions (Quirk, S. J., Ahkong, Q. F., Botham, G. M., Vos, J. & Lucy, J. A.) 159–167

Glycerol phosphate acyltransferase, adipose-tissue, epididymal, rat, inactivation of, catalysed by adenosine 3′:5′-cyclic monophosphate-dependent protein kinase and its possible involvement in the regulation of the biosynthesis of triacylglycerols (Nimmo, H. G. & Houston, B.) 607–610

Glycerolipids, effects of cortisol, corticotropin and thyroxine on the biosynthesis of, and on the activity of phosphatidate phosphatase in rat liver (Glenny, H. P. & Brindley, D. N.) 777–784

Glycine, inhibition by pentachlorophenol and related uncouplers of oxidative phosphorylation of the transport of proline and, by *Bacillus subtilis* (Nicholas, R. A. & Ordal, G. W.) 639–647

Glycine max, *see* Bean, soya

Glycine, oxidation of, by kidney-bean root-nodule leghaemoglobin and associated catabolic reactions at the haem group (Lehtovaara, P.) 351–358

Glycogen, effects of phenylephrine, vasopressin and glucagon on gluconeogenesis and on enzymes involved in the metabolism of, in isolated rat hepatocytes (Hue, L., Felíu, J. E. & Hers, H.-G.) 791–797

Glycogen, turnover of, in foetal rat liver (Gilbert, M. & Bourbon, J.) 785–789

Glycoproteins, fucosylated, plasma-membrane, neuroblastoma-cell, N2a, mouse, growth and the metabolism of (Milenkovic, A. G., Rachmeler, M. & Johnson, T. C.) 695–704

Glycoproteins, role of rat liver non-parenchymal cells in the clearance from plasma of, with terminal mannose and *N*-acetylglucosamine residues (Schlesinger, P. H., Doebber, T. W., Mandell, B. F., White, R., DeSchryver, C., Rodman, J. S., Miller, M. J. & Stahl, P.) 103–109

Gonadotropin, chorionic, human, induction by the β-subunit of, of the production of testosterone by mouse testis Leydig cells (Ramakrishnan, S., Das, C. & Talwar, G. P.) 599–602
Granulocytes, *see* Leucocytes, polymorphonuclear
Growth hormone, *see* Somatotropin
Growth, onset of the degradation of ribosomes during cessation of, in baby-hamster kidney BHK-21/C13 cells (Melvin, W. T. & Keir, H. M.) 933–941
Growth, turnover of protein in chick skeletal muscle during (Maruyama, K., Sunde, M. L. & Swick, R. W.) 573–582
Guanosine 3′:5′-cyclic monophosphate, method for the measurement of relative changes in the concentration of, in mouse neuroblastoma cells on muscarinic cholinergic stimulation (Strange, P. G.) 583–590
Guanosine 3′:5′-cyclic monophosphate, regulation of the biosynthesis of, in cultured mouse neuroblastoma cells (Bartfai, T., Breakefield, X. O. & Greengard, P.) 119–127
Guanosine [$\beta\gamma$-imido]triphosphate, effects of cholera toxin and, on the affinity of β-adrenergic receptors of toad erythrocyte membranes (Fischer, J. & Sharp, G. W. G.) 505–510

Haem group, oxidation of glycine by kidney-bean root-nodule leghaemoglobin and associated catabolic reactions at (Lehtovaara, P.) 351–358
Haem, kinetic evidence for the formation of a haem–oxygen complex during the degradation of, *in vitro* (Brown, S. B. & Thomas, S. E.) 327–330
Heart muscle, chicken, content and turnover of collagen in, and skeletal muscle during stretch-induced hypertrophy of skeletal muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 419–427
Heart muscle, chicken, rates of biosynthesis of protein in, and skeletal muscle and smooth muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 393–401
Heart muscle, role of phosphoenolpyruvate carboxykinase in the metabolism of amino acids in skeletal muscle and, of various animal species (Newsholme, E. A. & Williams, T.) 623–626
Heart, ox, thermodynamic analysis of the interaction between the mitochondrial coupling adenosine triphosphatase from, and its naturally occurring inhibitor protein (Gomez-Fernandez, J. C. & Harris, D. A.) 967–975
Heart, rat, anion/calcium ion ratios and production of protons during the uptake of calcium ions by mitochondria from, and liver (Harris, E. J.) 983–991
Heart, rat, perfused, steady-state, systematic variations in the concentrations of purine nucleotides in (Bates, D. J., Perrett, D. & Mowbray, J.) 485–493
Heart, rat, role of calcium ions in the regulation of the activity of mitochondrial nicotinamide–adenine dinucleotide-linked isocitrate dehydrogenase from, and other tissues (Denton, R. M., Richards, D. A. & Chin, J. G.) 899–906
Heart, rat, variation with age and sex of the concentrations of carnitine in, and other tissues (Borum, P. R.) 677–681
Hen, *see* Chicken
Hepatocytes, rabbit, isolated, effects of alloxan-induced diabetes on gluconeogenesis and ureogenesis in (Zaleski, J. & Bryła, J.) 563–568
Hepatocytes, rat and guinea-pig, isolated, differential effects of tryptophan on gluconeogenesis in (Smith, S. A., Elliott, K. R. F. & Pogson, C. I.) 817–825
Hepatocytes, rat, evidence for a reciprocal relationship between lipogenesis and ketogenesis in, from fed virgin and lactating animals (Benito, M. & Williamson, D. H.) 331–334
Hepatocytes, rat, isolated, characteristics of the transport of alanine, serine and glutamine across the plasma membrane of (Joseph, S. K., Bradford, N. M. & McGivan, J. D.) 827–836
Hepatocytes, rat, isolated, effect of the redox state of the cytosolic nicotinamide–adenine dinucleotide couple on the activity of adenosine 3′:5′-cyclic monophosphate phosphodiesterase and its role in the control of the responsiveness of, to glucagon (Clark, M. G. & Jarrett, I. G.) 805–816
Hepatocytes, rat, isolated, effects of phenylephrine, vasopressin and glucagon on gluconeogenesis and on enzymes involved in the metabolism of glycogen in (Hue, L., Feliu, J. E. & Hers, H.-G.) 791–797
Hepatocytes, rat, isolated, sources of ammonia for the biosynthesis of urea by (Krebs, H. A., Hems, R., Lund, P., Halliday, D. & Read, W. W. C.) 733–737
Hepatocytes, rat, isolated, stimulation by vasopressin, angiotensin and oxytocin of gluconeogenesis in (Whitton, P. D., Rodrigues, L. M. & Hems, D. A.) 893–898
Hepatoma cells, rat, cultured, effect of glutamine on the degradation of glutamine synthetase in, derived from Morris hepatoma 7288C (Crook, R. B. & Tomkins, G. M.) 47–52
Hepatoma cells, rat, cultured, extensive conjugation of metabolites of 3,4-dihydroxyphenethylamine in, and human skin fibroblasts (Crooks, P. A., Breakefield, X. O., Sulens, C. H., Castiglione, C. M. & Coward, J. K.) 187–196
Histidine, accelerated degradation of protein formed in rat and rabbit reticulocytes under conditions of depletion of, and other individual amino acids (Chandler, C. S. & Ballard, F. J.) 151–158
Hydrocortisone, *see* Cortisol
Hydrogen ions, anion/calcium ion ratios and production of, during the uptake of calcium ions by rat heart and liver mitochondria (Harris, E. J.) 983–991
Hydrolases, lysosomal, regulation of the biosynthesis of prostaglandins and of the selective release of, by mouse peritoneal macrophages (Bonney, R. J., Wightman, P. D., Davies, P., Sadowski, S. J., Kuehl, F. A., Jr. & Humes, J. L.) 433–442
3-Hydroxybutyrate, utilization of, for lipogenesis in lactating rat mammary gland *in vivo* (Robinson, A. M. & Williamson, D. H.) 635–638
1α-Hydroxy[7-^{3}H]cholecalciferol, synthesis of, and its metabolism in the chick (Edelstein, S., Noff, D., Freeman, D., Sheves, M. & Mazur, Y.) 111–117
3-Hydroxy-3-methylglutaryl-coenzyme A synthase, changes in the activities of acetoacetyl-coenzyme A deacylase and, in rat brain during postnatal development (Patel, (Patel, T. B. & Clark, J. B.) 951–958

5-Hydroxytryptamine, possible role of, as mediator in the action of calcitonin on rat gastrointestinal tract and pancreas (Nakhla, A. M. & Latif, A.) 339–342
Hyperglycaemia, induction of, and other metabolic effects of administration of 3-aminopicolinate to the rat (MacDonald, M. J., Huang, M.-T. & Lardy, H. A.) 495–504
Hypothalamus, rat, female, metabolism of high-molecular-weight ribonucleic acid in, and cerebral cortex during development (Hall, C. & Lim, L.) 511–521
Hypothalamus, sheep, binding of oestradiol-17β-receptor protein complex and of progesterone–receptor protein complex to chromatin from, from male and female animals (Perry, B. N. & Lopez, A.) 873–883

Inorganic sulphate, *see* Sulphate, inorganic
Insulin, effect of exogenous pyruvate on the induction by glucose of the release of, by isolated rat islets of Langerhans (Sener, A., Kawazu, S., Hutton, J. C., Boschero, A. C., Devis, G., Somers, G., Herchuelz, A. & Malaisse, W. J.) 217–232
Insulin, effects of, and other agents on the biosynthesis and release of amylase by rat parotid gland (McPherson, M. A. & Hales, C. N.) 855–863
Insulin, evidence for a role of, in the regulation of lipogenesis in lactating rat mammary gland (Robinson, A. M., Girard, J. R. & Williamson, D. H.) 343–346
Insulin, opposite effects of starvation on the oxidation of adenosine and on adenosine-induced release of, by isolated mouse islets of Langerhans (Andersson, A.) 619–621
Insulin, stimulation by pyruvate and lactate of the biosynthesis of proinsulin and the release of, by isolated rat islets of Langerhans (Jain, K., Asina, S. & Logothetopoulos, J.) 31–37
Intestine, small, rat, possible role of 5-hydroxytryptamine as mediator in the action of calcitonin on, and stomach and pancreas (Nakhla, A. M. & Latif, A.) 339–342
Islets of Langerhans, pancreatic, mouse, isolated, opposite effects of starvation on the oxidation of adenosine and on adenosine-induced release of insulin by (Andersson, A.) 619–621
Islets of Langerhans, pancreatic, rat, isolated, effect of exogenous pyruvate on the induction by glucose of the release of insulin by (Sener, A., Kawazu, S., Hutton, J. C., Boschero, A. C., Devis, G., Somers, G., Herchuelz, A. & Malaisse, W. J.) 217–232
Islets of Langerhans, pancreatic, rat, isolated, stimulation by pyruvate and lactate of the biosynthesis of proinsulin and the release of insulin by (Jain, K., Asina, S. & Logothetopoulos, J.) 31–37
Isocitrate dehydrogenase, nicotinamide–adenine dinucleotide-linked, role of calcium ions in the regulation of the activity of, in mitochondria from rat heart and other tissues (Denton, R. M., Richards, D. A. & Chin, J. G.) 899–906
Isocitrate lyase, effect of acetate on the rate of biosynthesis of, in *Chlorella fusca* var. *vacuolata* (Dunham, S. M. & Thurston, C. F.) 179–185
Isoleucyl-transfer ribonucleic acid synthetase, inhibition by pseudomonic acid of the activity of, in *Escherichia coli* M.R.E. 600 (Hughes, J. & Mellows, G.) 305–318
Isoproterenol, effects of, and other agents on the biosynthesis and release of amylase by rat parotid gland (McPherson, M. A. & Hales, C. N.) 855–863

Ketogenesis, evidence for a reciprocal relationship between lipogenesis and, in hepatocytes from fed virgin and lactating rats (Benito, M. & Williamson, D. H.) 331–334
Ketogenesis, regulation of, in the rat during postnatal development (Ferré, P., Pégorier, J.-P., Williamson, D. H. & Girard, J. R.) 759–765
Ketone bodies, regulation of the formation of, in the rat during postnatal development (Ferré, P., Pégorier, J.-P., Williamson, D. H. & Girard, J. R.) 759–765
Kidney bean, *see* Bean, kidney
Kidney cells (BHK-21/C13), baby-hamster, onset of the degradation of ribosomes during cessation of growth in (Melvin, W. T. & Keir, H. M.) 933–941
Kidney cortex, dog, interconversion of renin and high-molecular-weight renin in (Funakawa, S., Funae, Y. & Yamamoto, K.) 977–981
Krebs II ascites cells, characterization of the acidic ribosomal phosphoprotein L_7 from, and its relationship to *Escherichia coli* ribosomal proteins L7 and L12 (Leader, D. P. & Coia, A. A.) 569–572

Lactate, stimulation by pyruvate and, of the biosynthesis of proinsulin and the release of insulin by isolated rat islets of Langerhans (Jain, K., Asina, S. & Logothetopoulos, J.) 31–37
Lactation, characterization of lipid-linked glucose-containing oligosaccharides formed in rabbit mammary gland during (Speake, B. K. & White, D. A.) 993–1000
Lactation, evidence for a reciprocal relationship between lipogenesis and ketogenesis in hepatocytes from fed virgin rats and from fed rats during (Benito, M. & Williamson, D. H.) 331–334
Lactation, evidence for a role of insulin in the regulation of lipogenesis in rat mammary gland during (Robinson, A. M., Girard, J. R. & Williamson, D. H.) 343–346
Lactation, utilization of 3-hydroxybutyrate for lipogenesis in rat mammary gland *in vivo* during (Robinson, A. M. & Williamson, D. H.) 635–638
Lecithin, *see* Phosphatidylcholine
Leghaemoglobin, root-nodule, kidney-bean, oxidation of glycine by, and associated catabolic reactions at the haem group (Lehtovaara, P.) 351–358
Leghaemoglobins, root-nodule, leguminous-plant, formation of isomers of biliverdin during the degradation of, from various species (Lehtovaara, P. & Perttilä, U.) 359–364
Lens capsule, rat, biosynthesis and partial characterization of collagenous protein components of (Heathcote, J. G., Sear, C. H. J. & Grant, M. E.) 283–294
Leucine, effect of the administration of corticotropin on the incorporation of, into a specific cytosolic protein in rat adrenal gland (Dazord, A., Gallet, D. & Saez, J. M.) 233–239
Leucocytes, mononuclear, peritoneal, mouse, cultured, enhancement by double-stranded polyribonucleotides of the production by, of differentiation-stimulating factor(s) for mouse myeloid leukaemic cells (Tomida, M., Takenaga, K., Yamamoto, Y. & Hozumi, M.) 665–669

Leucocytes, mononuclear, peritoneal, mouse, regulation of the biosynthesis of prostaglandins and of the selective release of lysosomal hydrolases by (Bonney, R. J., Wightman, P. D., Davies, P., Sadowski, S. J., Kuehl, F. A., Jr. & Humes, J. L.) 433–442

Leucocytes, polymorphonuclear, human, subcellular localization of the activities of reduced nicotinamide–adenine dinucleotide (phosphate) oxidases in (Iverson, D. B., Wang-Iverson, P., Spitznagel, J. K. & DeChatelet, L. R.) 175–178

Leukaemic cells, myeloid, mouse, enhancement by double-stranded polyribonucleotides of the production by cultured mouse peritoneal mononuclear leucocytes of differentiation-stimulating factor(s) for (Tomida, M., Takenaga, K., Yamamoto, Y. & Hozumi, M.) 665–669

Leydig cells, testis, mouse, induction by the β-subunit of human chorionic gonadotropin of the production of testosterone by (Ramakrishnan, S., Das, C. & Talwar, G. P.) 599–602

Light, protective role of carotenoids in the sensitivity towards, of respiration by *Neurospora crassa* mitochondria (Ramadan-Talib, Z. & Prebble, J.) 767–775

Lipid, formation and characterization of glucose-containing oligosaccharides linked to, of lactating rabbit mammary gland (Speake, B. K. & White, D. A.) 993–1000

Lipogenesis, evidence for a reciprocal relationship between ketogenesis and, in hepatocytes from fed virgin and lactating rats (Benito, M. & Williamson, D. H.) 331–334

Lipogenesis, evidence for a role of insulin in the regulation of, in lactating rat mammary gland (Robinson, A. M., Girard, J. R. & Williamson, D. H.) 343–346

Lipogenesis, utilization of 3-hydroxybutyrate for, in lactating rat mammary gland *in vivo* (Robinson, A. M. & Williamson, D. H.) 635–638

Lipoprotein lipase, post-translational regulation of the activity of, in rat epididymal adipose tissue (Ashby, P., Bennett, D. P., Spencer, I. M. & Robinson, D. S.) 865–872

Lipoproteins, plasma, very-low-density, stimulation by, of the biosynthesis of triacylglycerols in isolated rat adipocytes (Thomopoulos, P., Berthelier, M., Lagrange, D., Chapman, M. J. & Laudat, M.-H.) 169–174

Liver cells, rabbit, isolated, effects of alloxan-induced diabetes on gluconeogenesis and ureogenesis in (Zaleski, J. & Bryła, J.) 563–568

Liver cells, rat and guinea-pig, isolated, differential effects of tryptophan on gluconeogenesis in (Smith, S. A., Elliot, K. R. F. & Pogson, C. I.) 817–825

Liver cells, rat, evidence for a reciprocal relationship between lipogenesis and ketogenesis in, from fed virgin and lactating animals (Benito, M. & Williamson, D. H.) 331–334

Liver cells, rat, isolated, characteristics of the transport of alanine, serine and glutamine across the plasma membrane of (Joseph, S. K., Bradford, N. M. & McGivan, J. D.) 827–836

Liver cells, rat, isolated, effect of the redox state of the cytosolic nicotinamide–adenine dinucleotide couple on the activity of adenosine 3′:5′-cyclic monophosphate phosphodiesterase and its role in the control of the responsiveness of, to glucagon (Clark, M. G. & Jarrett, I. G.) 805–816

Liver cells, rat, isolated, effects of phenylephrine, vasopressin and glucagon on gluconeogenesis and on enzymes involved in the metabolism of glycogen in (Hue, L., Felíu, J. E. & Hers, H.-G.) 791–797

Liver cells, rat, isolated, sources of ammonia for the biosynthesis of urea by (Krebs, H. A., Hems, R., Lund, P., Halliday, D. & Read, W. W. C.) 733–737

Liver cells, rat, isolated, stimulation by vasopressin, angiotensin and oxytocin of gluconeogenesis in (Whitton, P. D., Rodrigues, L. M. & Hems, D. A.) 893–898

Liver, chicken, utilization of methylmalonate for the biosynthesis of branched-chain fatty acids by preparations of (Scaife, J. R., Wahle, K. W. J. & Garton, G. A.) 799–804

Liver, rat, anion/calcium ion uptake and production of protons during the uptake of calcium ions by mitochondria from, and heart (Harris, E. J.) 983–991

Liver, rat, choice of precursors for the measurement of the turnover of proteins by a double-isotope method and its application to the study of mitochondrial proteins of (Burgess, R. J., Walker, J. H. & Mayer, R. J.) 919–926

Liver, rat, distribution of ^{14}C in glucose 6-phosphate biosynthesized from [1-^{14}C]ribose 5-phosphate by an extract from, and its relevance to the pentose phosphate pathway for the metabolism of glucose (Williams, J. F., Clark, M. G. & Blackmore, P. F.) 241–256

Liver, rat, effects of ammonium chloride and bicarbonate on the activity of glutaminase in mitochondria isolated from (Joseph, S. K. & McGivan, J. D.) 837–844

Liver, rat, effects of cortisol, corticotropin and thyroxine on the biosynthesis of glycerolipids and on the activity of phosphatidate phosphatase in (Glenny, H. P. & Brindley, D. N.) 777–784

Liver, rat, effects of glucagon and 6-*N*,2′-*O*-dibutyryladenosine 3′:5′-cyclic monophosphate on the transport of calcium ions by mitochondria isolated from (Hughes, B. P. & Barritt, G. J.) 295–304

Liver, rat, evidence for new reaction sequences in the pentose phosphate pathway for the metabolism of glucose by an extract of (Williams, J. F., Blackmore, P. F. & Clark, M. G.) 257–282

Liver, rat, evidence for the participation of aspartate aminotransferase in the formation of glucose in, of suckling newborn animals (Ferré, P. & Williamson, D. H.) 335–338

Liver, rat, evidence for the presence of a subpopulation of membrane-bound ribosomes from, that are detached *in vitro* by carcinogens and by centrifugation (Palmer, D. N., Rabin, B. R. & Williams, D. J.) 9–14

Liver, rat, evidence from essential-fatty acid-deficient animals for a regulatory role of choline kinase in the biosynthesis of phosphatidylcholine in (Infante, J. P. & Kinsella, J. E.) 631–633

Liver, rat, evidence that most of the membrane proteins of lysosomes of, are exposed to the cytoplasm (Schneider, D. L., Burnside, J., Gorga, F. R. & Nettleton, C. J.) 75–82

Liver, rat, foetal, turnover of glycogen in (Gilbert, M. & Bourbon, J.) 785–789
Liver, rat, hydrolysis of phosphatidylinositol by lysosomal enzymes of, and brain (Irvine, R. F., Hemington, N. & Dawson, R. M. C.) 475–484
Liver, rat, influence of respiratory state on the activity of monoamine oxidase in mitochondria isolated from (Smith, G. S. & Reid, R. A.) 1011–1014
Liver, rat, initiation of the transcription of deoxyribonucleic acid for the biosynthesis of ribonucleic acid in normal and phosphorylated nucleosomes from (Ord, M. G. & Stocken, L. A.) 615–618
Liver, rat, location of the activity of coproporphyrinogen III oxidase in mitochondria from (Grandchamp, B., Phung, N. & Nordmann, Y.) 97–102
Liver, rat, mechanisms of the effects of the administration of ethanol and glycerol on the activity of phosphatidate phosphatase in (Savolainen, M. J. & Hassinen, I. E.) 885–892
Liver, rat, participation of cysteine and cystine in the inactivation of tyrosine aminotransferase in homogenates of (Buckley, W. T. & Milligan, L. P.) 449–454
Liver, rat, perfused, isolated, activity of phenol sulphotransferase and the metabolism of inorganic sulphate in (Mulder, G. J. & Keulemans, K.) 959–965
Liver, rat, perfused, stimulation by vasopressin of the activity of pyruvate dehydrogenase in (Hems, D. A., McCormack, J. G. & Denton, R. M.) 627–629
Liver, rat, regulation of the extramitochondrial concentration of calcium ions by mitochondria from (Nicholls, D. G.) 463–474
Liver, rat, relative rates of turnover of subunits of mitochondrial proteins of (Walker, J. H., Burgess, R. J. & Mayer, R. J.) 927–932
Liver, rat, role of non-parenchymal cells of, in the clearance from plasma of glycoproteins with terminal mannose and *N*-acetylglucosamine residues (Schlesinger, P. H., Doebber, T. W., Mandell, B. F., White, R., DeSchryver, C., Rodman, J. S., Miller, M. J. & Stahl, P.) 103–109
Liver, rat, role of reduced glutathione in the reduction of diamide by mitochondria isolated from (Jocelyn, P. C.) 649–664
Liver, rat, stable enhancement of the retention of calcium ions by mitochondria isolated from, after the administration of glucagon to the animal (Prpić, V., Spencer, T. L. & Bygrave, F. L.) 705–714
Liver, rat, use of rat liver nucleoplasm for the characterization of the biosynthesis of heterogeneous nuclear ribonucleic acid *in vivo* (Beebee, T. J. C.) 715–725
Liver, rat, variation with age and sex of the concentrations of carnitine in, and other tissues (Borum, P. R.) 677–681
Lung, rabbit, accumulation and turnover of pulmonary surfactant in (Desai, R., Tetley, T. D., Curtis, C. G., Powell, G. M. & Richards, R. J.) 455–462
Lysosomes, fibroblast, effect of inhibitors of endocytosis on the intracellular and extracellular activities of enzymes of (von Figura, K. & Weber, E.) 943–950
Lysosomes, liver and brain, rat, hydrolysis of phosphatidylinositol by enzymes of (Irvine, R. F., Hemington, N. & Dawson, R. M. C.) 475–484
Lysosomes, liver, rat, evidence that most of the membrane proteins of, are exposed to the cytoplasm (Schneider, D. L., Burnside, J., Gorga, F. R. & Nettleton, C. J.) 75–82

Macrophages, peritoneal, mouse, regulation of the biosynthesis of prostaglandins and of the selective release of lysosomal hydrolases by (Bonney, R. J., Wightman, P. D., Davies, P., Sadowski, S. J., Kuehl, F. A., Jr. & Humes, J. L.) 433–442
Macrophages, *see also* Leucocytes, mononuclear
Malondialdehyde, evidence that the formation of, in human platelets is catalysed by thromboxane synthase (McMillan, R. M., MacIntyre, D. E., Booth, A. & Gordon, J. L.) 595–598
Mammary-gland adenocarcinoma R3230AC cells, rat, sodium ion-dependent and -independent transport of 2-aminoisobutyrate into, and the effects of streptozotocin-induced diabetes in the host animal (Hissin, P. J. & Hilf, R.) 205–215
Mammary gland, rabbit, lactating, formation and characterization of lipid-linked glucose-containing oligosaccharides of (Speake, B. K. & White, D. A.) 993–1000
Mammary gland, rat, lactating, evidence for a role of insulin in the regulation of lipogenesis in (Robinson, A. M., Girard, J. R. & Williamson, D. H.) 343–346
Mammary gland, rat, lactating, utilization of 3-hydroxybutyrate for lipogenesis in, *in vivo* (Robinson, A. M. & Williamson, D. H.) 635–638
Mannose residues, terminal, role of rat liver non-parenchymal cells in the clearance from plasma of glycoproteins with, and *N*-acetylglucosamine residues (Schlesinger, P. H., Doebber, T. W., Mandell, B. F., White, R., DeSchryver, C., Rodman, J. S., Miller, M. J. & Stahl, P.) 103–109
Membrane, chorioallantoic, chick-embryo, functional involvement of carbonic anhydrase in the transport of calcium ions by (Tuan, R. S. & Zrike, J.) 67–74
Membrane, microsomal, liver, rat, evidence for the presence of a subpopulation of ribosomes bound to, that are detached *in vitro* by carcinogens and by centrifugation (Palmer, D. N., Rabin, B. R. & Williams, D. J.) 9–14
Membrane, plasma, characteristics of the transport of alanine, serine and glutamine across, of isolated rat hepatocytes (Joseph, S. K., Bradford, N. M. & McGivan, J. D.) 827–836
Membrane, plasma, neuroblastoma-cell, N2a, mouse, growth and the metabolism of fucosylated glycoproteins of (Milenkovic, A. G., Rachmeler, M. & Johnson, T. C.) 695–704
Membrane proteins, erythrocyte, human, characterization of, after cell fusion induced by oleoylglycerol in the presence of calcium ions (Quirk, S. J., Ahkong, Q. F., Botham, G. M., Vos, J. & Lucy, J. A.) 159–167
Membrane proteins, lysosomal, liver, rat, evidence that most of, are exposed to the cytoplasm (Schneider, D. L., Burnside, J., Gorga, F. R. & Nettleton, C. J.) 75–82
Membranes, adipocyte, epididymal, rat, modulation by thyroid hormones of the numbers of β-adrenergic receptors in (Giudicelli, Y.) 1007–1010

Membranes, erythrocyte, toad, effects of cholera toxin and guanosine [$\beta\gamma$-imido]triphosphate on the affinity of the β-adrenergic receptors of (Fischer, J. & Sharp, G. W. G.) 505–510
Mersalyl, chemical-modification approach to elucidation of the olfactory code exemplified by the inhibition by, of the response of frog olfactory mucosa to pentyl acetate and related odorants (Menevse, A., Dodd, G. & Poynder, T. M.) 845–854
3-Methoxytyramine 4(3)-*O*-glucuronide, identification of, as a major metabolite of 3,4-dihydroxyphenethylamine in cultured rat hepatoma cells (Crooks, P. A., Breakefield, X. O., Sulens, C. H., Castiglione, C. M. & Coward, J. K.) 187–196
3-Methoxytyramine 4-*O*-sulphate, identification of, as a major metabolite of 3,4-dihydroxyphenethylamine in cultured human skin fibroblasts (Crooks, P. A., Breakefield, X. O., Sulens, C. H., Castiglione, C. M. & Coward, J. K.) 187–196
Methylmalonate, utilization of, for the biosynthesis of branched-chain fatty acids by preparations of chicken liver and sheep inguinal adipose tissue (Scaife, J. R., Wahle, K. W. J. & Garton, G. A.) 799–804
Microsomal fraction, pancreatic, rat, sensitivity towards proteinases of the release of α-amylase and ribonuclease by (Pearce, P. D., May, B. K. & Elliott, W. H.) 611–614
Mitochondria, brain, rat, changes in the activities of acetoacetyl-coenzyme A deacylase and 3-hydroxy-3-methylglutaryl-coenzyme A synthase in, during postnatal development (Patel, T. B. & Clark, J. B.) 951–958
Mitochondria, heart and liver, rat, anion/calcium ion ratios and production of protons during the uptake of calcium ions by (Harris, E. J.) 983–991
Mitochondria, heart, ox, thermodynamic analysis of the interaction between the coupling adenosine triphosphatase from, and its naturally occurring inhibitor protein (Gomez-Fernandez, J. C. & Harris, D. A.) 967–975
Mitochondria, heart, rat, role of calcium ions in the regulation of the activity of nicotinamide–adenine dinucleotide-linked isocitrate dehydrogenase of, and of mitochondria from other rat tissues (Denton, R. M., Richards, D. A. & Chin, J. G.) 899–906
Mitochondria, liver, rat, choice of precursors for the measurement of the turnover of proteins by a double-isotope method and its application to the study of proteins of (Burgess, R. J., Walker, J. H. & Mayer, R. J.) 919–926
Mitochondria, liver, rat, isolated, effects of ammonium chloride and bicarbonate on the activity of glutaminase in (Joseph, S. K. & McGivan, J. D.) 837–844
Mitochondria, liver, rat, isolated, effects of glucagon and 6-*N*,2′-*O*-dibutyryladenosine 3′:5′-cyclic monophosphate on the transport of calcium ions by (Hughes, B. P. & Barritt, G. J.) 295–304
Mitochondria, liver, rat, isolated, enhancement of the retention of calcium ions by, after the administration of glucagon to the animal (Prpić, V., Spencer, T. L. & Bygrave, F. L.) 705–714
Mitochondria, liver, rat, isolated, influence of respiratory state on the activity of monoamine oxidase in (Smith, G. S. & Reid, R. A.) 1011–1014
Mitochondria, liver, rat, isolated, role of reduced glutathione in the reduction of diamide by (Jocelyn, P. C.) 649–664
Mitochondria, liver, rat, location of the activity of coproporphyrinogen III oxidase in (Grandchamp, B., Phung, N. & Nordmann, Y.) 97–102
Mitochondria, liver, rat, regulation of the extramitochondrial concentration of calcium ions by (Nicholls, D. G.) 463–474
Mitochondria, liver, rat, relative rates of turnover of subunits of proteins of (Walker, J. H., Burgess, R. J. & Mayer, R. J.) 927–932
Mitochondria, mung-bean, cyanide-insensitive uptake of oxygen by, in the presence of ascorbate/*NNN′N′*-tetramethyl-*p*-phenylenediamine mixture as electron donor (Wilson, S. B.) 129–136
Mitochondria, *Neurospora crassa*, protective role of carotenoids in the photosensitivity of respiration by (Ramadan-Talib, Z. & Prebble, J.) 767–775
Modeccin, inhibitory effect of, on the elongation of polypeptide chains during the biosynthesis of protein catalysed by brine-shrimp ribosomes (Montanaro, L., Sperti, S., Zamboni, M., Denaro, M., Testoni, G., Gasperi-Campani, A. & Stirpe, F.) 371–379
Monoamine oxidase, influence of respiratory state on the activity of, in isolated rat liver mitochondria (Smith, G. S. & Reid, R. A.) 1011–1014
Morris hepatoma 7288C, rat, effect of glutamine on the degradation of glutamine synthetase in cultured hepatoma cells derived from (Crook, R. B. & Tomkins, G. M.) 47–52
Müllerian ducts, chick-embryo, differentiating, effects of oestrogens on the activity of ornithine decarboxylase and the concentrations of polyamines in, and other organs (Teng, C. S. & Teng, C. T.) 143–149
Mung bean, *see* Bean, mung
Muscle, heart, chicken, content and turnover of collagen in, and skeletal muscle during stretch-induced hypertrophy of skeletal muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 419–427
Muscle, heart, chicken, rates of biosynthesis of protein in, and skeletal muscle and smooth muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 393–401
Muscle, heart, role of phosphoenolpyruvate carboxykinase in the metabolism of amino acids in skeletal muscle and, of various animal species (Newsholme, E. A. & Williams, T.) 623–626
Muscle, skeletal, chick, turnover of protein in, during growth (Maruyama, K., Sunde, M. L. & Swick, R. W.) 573–582
Muscle, skeletal, chicken, changes in the rates of biosynthesis and degradation of protein in, during stretch-induced hypertrophy of skeletal muscle (Laurent, G. J., Sparrow, M. P. & Millward, D. J.) 407–417
Muscle, skeletal, chicken, content and turnover of collagen in, and heart muscle during stretch-induced hypertrophy of skeletal muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 419–427
Muscle, skeletal, chicken, rates of biosynthesis of protein in, and heart muscle and smooth muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 393–401

Muscle, skeletal, mouse, composition and metabolism of phospholipids in, and nervous tissue of normal animals and of animals with muscular dystrophy (Kwok, C. T. & Austin, L.) 15–22
Muscle, skeletal, mouse, purification and properties of a factor inhibitory towards the biosynthesis of protein from, of normal animals and those with muscular dystrophy (Petryshyn, R. A. & Nicholls, D. M.) 907–917
Muscle, skeletal, rat, active and immobilized, effects of starvation on the turnover of protein and the concentration of ribonucleic acid in (Goldspink, D. F.) 603–606
Muscle, skeletal, rat, effect of surgical trauma on the turnover of protein in (Hoover-Plow, J. L. & Clifford, A. J.) 137–142
Muscle, skeletal, rat, regulation by adrenaline of the activity of phosphorylase phosphatase in (Tao, S.-H., Huang, F. L., Lynch, A. & Glinsmann, W. H.) 347–350
Muscle, skeletal, rat, variation with age and sex of the concentrations of carnitine in, and other tissues (Borum, P. R.) 677–681
Muscle, skeletal, role of phosphoenolpyruvate carboxykinase in the metabolism of amino acids in heart muscle and, of various animal species (Newsholme, E. A. & Williams, T.) 623–626
Muscle, smooth, chicken, rates of biosynthesis of protein in, and skeletal muscle and heart muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 393–401
Muscular dystrophy, composition and metabolism of phospholipids in nervous tissue and skeletal muscle of normal mice and of mice with (Kwok, C. T. & Austin, L.) 15–22
Muscular dystrophy, purification and properties of a factor inhibitory towards the biosynthesis of protein from skeletal muscle of normal mice and those with (Petryshyn, R. A. & Nicholls, D. M.) 907–917
Myofibrils, skeletal-muscle, rat, effect of surgical trauma on the turnover of protein in, and sarcoplasm (Hoover-Plow, J. L. & Clifford, A. J.) 137–142

Nervous tissue, mouse, composition and metabolism of phospholipids in, and skeletal muscle of normal animals and of animals with muscular dystrophy (Kwok, C. T. & Austin, L.) 15–22
Neuroblastoma cells, mouse, cultured, regulation of the biosynthesis of guanosine 3′:5′-cyclic monophosphate in (Bartfai, T., Breakefield, X. O. & Greengard, P.) 119–127
Neuroblastoma cells, mouse, method for the measurement of relative changes in the concentration of guanosine 3′:5′-cyclic monophosphate in, on muscarinic cholinergic stimulation (Strange, P. G.) 583–590
Neuroblastoma cells, N2a, mouse, growth and the metabolism of plasma-membrane fucosylated glycoproteins in (Milenkovic, A. G., Rachmeler, M. & Johnson, T. C.) 695–704
Neurospora crassa, protective role of carotenoids in the photosensitivity of respiration by mitochondria from (Ramadan-Talib, Z. & Prebble, J.) 767–775
Nicotinamide–adenine dinucleotide couple, cytosolic, effect of the redox state of, on the activity of adenosine 3′:5′-cyclic monophosphate phosphodiesterase and its role in the control of the responsiveness of isolated rat hepatocytes to glucagon (Clark, M. G. & Jarrett, I. G.) 805–816
Nicotinamide–adenine dinucleotide (phosphate) (reduced) oxidases, subcellular localization of the activities of, in human polymorphonuclear leucocytes (Iverson, D. B., Wang-Iverson, P., Spitznagel, J. K. & DeChatelet, L. R.) 175–178
Non-histone protein, *see* Protein, non-histone
sym-Nor-spermine (11,11-diamino-4,8-diazaundecane), pathway for the biosynthesis of, and other polyamines in *Caldariella acidophila* (De Rosa, M., De Rosa, S., Gambacorta, A., Carteni-Farina, M. & Zappia, V.) 1–7
Nuclear ribonucleic acid, *see* Ribonucleic acid, nuclear
Nuclei, hypothalamus, sheep, binding of oestradiol-17β-receptor protein complex and of progesterone-receptor protein complex to chromatin from, from male and female animals (Perry, B. N. & Lopez, A.) 873–883
Nuclei, uterus, rat, binding characteristics, content and distribution of oestrogen-receptor proteins in, and cytosol in relation to intra-uterine devices and the oestrous cycle (Myatt, L., Chaudhuri, G., Elder, M. G. & Lim, L.) 523–529
Nuclei, ventral-prostate-gland, rat, effects of polyamines on the reactions catalysed by protein kinases associated with chromatin and non-histone proteins of (Ahmed, K., Wilson, M. J., Goueli, S. A. & Williams-Ashman, H. G.) 739–750
Nucleoplasm, liver, rat, use of, for the characterization of the biosynthesis of heterogeneous nuclear ribonucleic acid in rat liver *in vivo* (Beebee, T. J. C.) 715–725
Nucleosomes, liver, rat, normal and phosphorylated, initiation of the transcription of deoxyribonucleic acid for the biosynthesis of ribonucleic acid in (Ord, M. G. & Stocken, L. A.) 615–618

Obelin, use of, in a study of the inhibition by calcium ions of the formation of adenosine 3′:5′-cyclic monophosphate in sealed pigeon erythrocyte 'ghosts' (Campbell, A. K. & Dormer, R. L.) 53–66
Oestradiol-17β, effects of, on the activity of ornithine decarboxylase and the concentrations of polyamines in differentiating chick-embryo Müllerian ducts and other organs (Teng, C. S. & Teng, C. T.) 143–149
Oestradiol-17β-receptor protein complex, binding of progesterone-receptor protein complex and, to hypothalamus chromatin from male and female sheep (Perry, B. N. & Lopez, A.) 873–883
Oestrogen-receptor proteins, uterus, rat, binding characteristics, content and intracellular distribution of, in relation to intra-uterine devices and the oestrous cycle (Myatt, L., Chaudhuri, G., Elder, M. G. & Lim, L.) 523–529
Oestrogens, effects of, on the activity of ornithine decarboxylase and the concentrations of polyamines in differentiating chick-embryo Müllerian ducts and other organs (Teng, C. S. & Teng, C. T.) 143–149

Oestrous cycle, binding characteristics, content and intracellular distribution of oestrogen-receptor proteins in rat uterus in relation to intra-uterine devices and (Myatt, L., Chaudhuri, G., Elder, M. G. & Lim, L.) 523–529

Oleoylglycerol, characterization of human erythrocyte membrane proteins after cell fusion induced by, in the presence of calcium ions (Quirk, S. J., Ahkong, Q. F., Botham, G. M., Vos, J. & Lucy, J. A.) 159–167

Olfactory code, chemical-modification approach to elucidation of, exemplified by the inhibition by mersalyl of the response of frog olfactory mucosa to pentyl acetate and related odorants (Menevse, A., Dodd, G. & Poynder, T. M.) 845–854

Olfactory mucosa, frog, chemical-modification approach to elucidation of the olfactory code exemplified by the inhibition by mersalyl of the response of, to pentyl acetate and related odorants (Menevse, A., Dodd, G. & Poynder, T. M.) 845–854

Oligosaccharides, glucose-containing, lipid-linked, formation and characterization of, of lactating rabbit mammary gland (Speake, B. K. & White, D. A.) 993–1000

Ornithine decarboxylase, effects of oestrogens on the activity of, and the concentrations of polyamines in differentiating chick-embryo Müllerian ducts and other organs (Teng, C. S. & Teng, C. T.) 143–149

Ovalbumin, digestion of, by human pepsins 1, 3 and 5 (Walker, V. & Taylor, W. H.) 429–432

Oviduct, chicken, isolation of a high-molecular-weight high-mobility-group-type non-histone protein from (Teng, C. S., Gallagher, K. & Teng, C. T.) 1003–1006

Oxidative phosphorylation, *see* Phosphorylation, oxidative

Oxygen, cyanide-insensitive uptake of, by mung-bean mitochondria in the presence of ascorbate/*NNN'N'*-tetramethyl-*p*-phenylenediamine mixture as electron donor (Wilson, S. B.) 129–136

Oxygen, kinetic evidence for the formation of a complex between haem and, during the degradation of haem *in vitro* (Brown, S. B. & Thomas, S. E.) 327–330

Oxytocin, stimulation by vasopressin, angiotensin and, of gluconeogenesis in isolated rat hepatocytes (Whitton, P. D., Rodrigues, L. M. & Hems, D. A.) 893–898

Pancreas, mouse, opposite effects of starvation on the oxidation of adenosine and on adenosine-induced release of insulin by islets of Langerhans isolated from (Andersson, A.) 619–621

Pancreas, rat, effect of exogenous pyruvate on the induction by glucose of the release of insulin by islets of Langerhans isolated from (Sener, A., Kawazu, S., Hutton, J. C., Boschero, A. C., Devis, G., Somers, G., Herchuelz, A. & Malaisse, W. J.) 217–232

Pancreas, rat, possible role of 5-hydroxytryptamine as mediator in the action of calcitonin on, and gastrointestinal tract (Nakhla, A. M. & Latif, A.) 339–342

Pancreas, rat, sensitivity towards proteinases of the release of α-amylase and ribonuclease by microsomal fraction from (Pearce, P. D., May, B. K. & Elliott, W. H.) 611–614

Pancreas, rat, stimulation by pyruvate and lactate of the biosynthesis of proinsulin and the release of insulin by islets of Langerhans isolated from (Jain, K., Asina, S. & Logothetopoulos, J.) 31–37

Parotid gland, rat, control of the biosynthesis and release of amylase by (McPherson, M. A. & Hales, C. N.) 855–863

Parotid gland, rat, dependence on calcium ions of the inhibition by α-adrenergic agonists and cholinergic agonists of the biosynthesis of protein in slices of (Kanagasuntheram, P. & Lim, S. C.) 23–29

Pea (*Pisum sativum*), formation of isomers of biliverdin during the degradation of leghaemoglobins from root nodules of, and other leguminous plants (Lehtovaara, P. & Perttilä, U.) 359–364

Pentachlorophenol, inhibition by, and related uncouplers of oxidative phosphorylation of the transport of proline and glycine by *Bacillus subtilis* (Nicholas, R. A. & Ordal, G. W.) 639–647

Pentose phosphate pathway, distribution of ^{14}C in glucose 6-phosphate biosynthesized from [1-^{14}C]ribose 5-phosphate by a rat liver extract and its relevance to, for the metabolism of glucose (Williams, J. F., Clark, M. G. & Blackmore, P. F.) 241–256

Pentose phosphate pathway, evidence for new reaction sequences in, for the metabolism of glucose by a rat liver extract (Williams, J. F., Blackmore, P. F. & Clark, M. G.) 257–282

Pentyl acetate, chemical-modification approach to elucidation of the olfactory code exemplified by the inhibition by mersalyl of the response of frog olfactory mucosa to, and related odorants (Menevse, A., Dodd, G. & Poynder, T. M.) 845–854

Pepsins 1, 3 and 5, human, digestion of ovalbumin by (Walker, V. & Taylor, W. H.) 429–432

Phaseolus aureus, *see* Bean, mung

Phaseolus vulgaris, *see* Bean, kidney

Phenol sulphotransferase, activity of, and the metabolism of inorganic sulphate in isolated perfused rat liver (Mulder, G. J. & Keulemans, K.) 959–965

Phenylalanine, dependence on calcium ions of the inhibition by α-adrenergic agonists and cholinergic agonists of the incorporation of, into protein in rat parotid-gland slices (Kanagasuntheram, P. & Lim, S. C.) 23–29

Phenylephrine, effects of vasopressin, glucagon and, on gluconeogenesis and on enzymes involved in the metabolism of glycogen in isolated rat hepatocytes (Hue, L., Felíu, J. E. & Hers, H.-G.) 791–797

Phosphatidate phosphatase, effects of cortisol, corticotropin and thyroxine on the biosynthesis of glycerolipids and on the activity of, in rat liver (Glenny, H. P. & Brindley, D. N.) 777–784

Phosphatidate phosphatase, mechanisms of the effects of the administration of ethanol and glycerol on the activity of, in rat liver (Savolainen, M. J. & Hassinen, I. E.) 885–892

Phosphatidylcholine, concentration and metabolism of, and other phospholipids in nervous tissue and skeletal muscle of normal mice and of mice with muscular dystrophy (Kwok, C. T. & Austin, L.) 15–22

Phosphatidylcholine, evidence from essential-fatty acid-deficient animals for a regulatory role of choline kinase in the biosynthesis of, in rat liver (Infante, J. P. & Kinsella, J. E.) 631–633
Phosphatidylinositol, hydrolysis of, by lysosomal enzymes of rat liver and brain (Irvine, R. F., Hemington, N. & Dawson, R. M. C.) 475–484
Phosphatidyl-*myo*-inositol 4,5-bjsphosphate (triphosphoinositide), activation by calcium ions of the hydrolysis of phosphatidyl-*myo*-inositol 4-phosphate (diphosphoinositide) and, by guinea-pig brain synaptosomes (Griffin, H. D. & Hawthorne, J. N.) 541–552
Phosphatidyl-*myo*-inositol 4-phosphate (diphosphoinositide), activation by calcium ions of the hydrolysis of phosphatidyl-*myo*-inositol 4,5-bisphosphate (triphosphoinositide) and, by guinea-pig brain synaptosomes (Griffin, H. D. & Hawthorne, J. N.) 541–552
Phosphodiesterases, properties of the calcium ion-dependent protein modulator of the activities of, of mouse epidermis (Murray, A. W. & Rogers, A.) 727–732
Phosphoenolpyruvate carboxykinase, role of, in the metabolism of amino acids in skeletal muscle and heart muscle of various animal species (Newsholme, E. A. & Williams, T.) 623–626
Phospholipase A, stimulation by, of the content of prostaglandin E and the biosynthesis of somatotropin in rat anterior pituitary gland (Betteridge, A. & Wallis, M.) 319–323
Phospholipids, composition and metabolism of, in nervous tissue and skeletal muscle of normal mice and of mice with muscular dystrophy (Kwok, C. T. & Austin, L.) 15–22
Phosphoprotein L_{γ}, ribosomal, Krebs II-ascites-cell, characterization of, and its relationship to *Escherichia coli* ribosomal proteins L7 and L12 (Leader, D. P. & Coia, A. A.) 569–572
Phosphorylase phosphatase, regulation by adrenaline of the activity of, in rat skeletal muscle (Tao, S.-H., Huang, F. L., Lynch, A. & Glinsmann, W. H.) 347–350
Phosphorylation, oxidative, inhibition by pentachlorophenol and related uncouplers of, of the transport of proline and glycine by *Bacillus subtilis* (Nicholas, R. A. & Ordal, G. W.) 639–647
Pigeon, use of obelin in a study of the inhibition by calcium ions of the formation of adenosine 3′:5′-cyclic monophosphate in sealed erythrocyte 'ghosts' from (Campbell A. K. & Dormer, R. L.) 53–66
Pisum sativum, *see* Pea
Pituitary gland, anterior, rat, stimulation by phospholipase A of the content of prostaglandin E and the biosynthesis of somatotropin in (Betteridge, A. & Wallis, M.) 319–323
Plasma lipoproteins, *see* Lipoproteins, plasma
Plasma membrane, *see* Membrane, plasma
Plasma, rat, role of the liver non-parenchymal cells in the clearance from, of glycoproteins with terminal mannose and *N*-acetylglucosamine residues (Schlesinger, P. H., Doebber, T. W., Mandell, B. F., White, R., DeSchryver, C., Rodman, J. S., Miller, M. J. & Stahl, P.) 103–109
Plasma, rat, variation with age and sex of the concentrations of carnitine in, and other tissues (Borum, P. R.) 677–681
Platelets, human, broken and intact, inhibition by adenosine analogues of the activity of adenylate cyclase in preparations of (Haslam, R. J., Davidson, M. M. L. & Desjardins, J. V.) 83–95
Platelets, human, evidence that the formation of malondialdehyde in, is catalysed by thromboxane synthase (McMillan, R. M., MacIntyre, D. E., Booth, A. & Gordon, J. L.) 595–598
Polyamines, effect of, on the reactions catalysed by protein kinases associated with rat ventral-prostate-gland chromatin and non-histone proteins (Ahmed, K., Wilson, M. J., Goueli, S. A. & Williams-Ashman, H. G.) 739–750
Polyamines, effects of oestrogens on the activity of ornithine decarboxylase and the concentrations of, in differentiating chick-embryo Müllerian ducts and other organs (Teng, C. S. & Teng, C. T.) 143–149
Polyamines, pathway for the biosynthesis of, in *Caldariella acidophila* (De Rosa, M., De Rosa, S., Gambacorta, A., Carteni-Farina, M. & Zappia, V.) 1–7
Polymorphonuclear leucocytes, *see* Leucocytes, polymorphonuclear
Polypeptide chains, inhibitory effect of modeccin on the elongation of, during the biosynthesis of protein catalysed by brine-shrimp ribosomes (Montanaro, L., Sperti, S., Zamboni, M., Denaro, M., Testoni, G., Gasperi-Campani, A. & Stirpe, F.) 371–379
Polyribonucleotides, double-stranded, enhancement by, of the production by cultured mouse peritoneal mononuclear leucocytes of differentiation-stimulating factor(s) for mouse myeloid leukaemic cells (Tomida, M., Takenaga, K., Yamamoto, Y. & Hozumi, M.) 665–669
Progesterone-receptor protein complex, binding of oestradiol-17β-receptor protein complex and, to hypothalamus chromatin from male and female sheep (Perry, B. N. & Lopez, A.) 873–883
Progesterone-receptor protein, cytosolic, uterus, human, interaction of progestins with (Kasid, A., Buckshee, K., Hingorani, V. & Laumas, K. R.) 531–539
Progesterone, specific 11α-hydroxylation of, by *Aspergillus ochraceus* TS (Samanta, T. B., Roy, N. & Chattopadhyay, S.) 593–594
Progestins, interaction of, with the cytosolic progesterone-receptor protein of (Kasid, A., Buckshee, K., Hingorani, V. & Laumas, K. R.) 531–539
Proinsulin, stimulation by pyruvate and lactate of the biosynthesis of, and the release of insulin by isolated rat islets of Langerhans (Jain, K., Asina, S. & Logothetopoulos, J.) 31–37
Proline, inhibition by pentachlorophenol and related uncouplers of oxidative phosphorylation of the transport of glycine and, by *Bacillus subtilis* (Nicholas, R. A. & Ordal, G. W.) 639–647
Propionate, metabolism of, in relation to the biosynthesis of branched-chain fatty acids by preparations of chicken liver and sheep inguinal adipose tissue (Scaife, J. R., Wahle, K. W. J. & Garton, G. A.) 799–804
Prostaglandin E, stimulation by phospholipase A of the content of, and the biosynthesis of somatotropin in rat anterior pituitary gland (Betteridge, A. & Wallis, M.) 319–323

Prostaglandins, regulation of the biosynthesis of, and of the selective release of lysosomal hydrolases by mouse peritoneal macrophages (Bonney, R. J., Wightman, P. D., Davies, P., Sadowski, S. J., Kuehl, F. A., Jr. & Humes, J. L.) 433–442

Prostate gland, ventral, rat, effects of polyamines on the reactions catalysed by protein kinases associated with nuclear chromatin and non-histone proteins of (Ahmed, K., Wilson, M. J., Goueli, S. A. & Williams-Ashman, H. G.) 739–750

Protease, *see* Proteinase

Protein, accelerated degradation of, formed in rat and rabbit reticulocytes under conditions of depletion of individual amino acids (Chandler, C. S. & Ballard, F. J.) 151–158

Protein, adenosine triphosphatase-inhibitor, mitochondrial, heart, ox, thermodynamic analysis of the interaction between ox heart mitochondrial coupling adenosine triphosphatase and (Gomez-Fernandez, J. C. & Harris, D. A.) 967–975

Protein, analysis of errors in estimation of the rate of biosynthesis of, by constant infusion of a labelled amino acid into an animal (Garlick, P. J.) 402–405

Protein, changes in the rates of biosynthesis and degradation of, in chicken skeletal muscle during stretch-induced hypertrophy of skeletal muscle (Laurent, G. J., Sparrow, M. P. & Millward, D. J.) 407–417

Protein, cytosolic, specific, effect of the administration of corticotropin on the biosynthesis of, in rat adrenal gland (Dazord, A., Gallet, D. & Saez, J. M.) 233–239

Protein, dependence on calcium ions of the inhibition by α-adrenergic agonists and cholinergic agonists of the biosynthesis of, in rat parotid-gland slices (Kanagasuntheram, P. & Lim, S. C.) 23–29

Protein, effect of surgical trauma on the turnover of, in rat skeletal muscle (Hoover-Plow, J. L. & Clifford, A. J.) 137–142

Protein, effects of starvation on the concentration of ribonucleic acid and the turnover of, in active and immobilized rat skeletal muscle (Goldspink, D. F.) 603–606

Protein, inhibitory effect of modeccin on the elongation of polypeptide chains during the biosynthesis of, catalysed by brine-shrimp ribosomes (Montanaro, L., Sperti, S., Zamboni, M., Denaro, M., Testoni, G., Gasperi-Campani, A. & Stirpe, F.) 371–379

Protein kinase, adenosine 3′:5′-cyclic monophosphate-dependent, inactivation of rat epididymal adipose-tissue glycerol phosphate acyltransferase catalysed by, and its possible involvement in the regulation of the biosynthesis of triacylglycerols (Nimmo, H. G. & Houston, B.) 607–610

Protein kinases, effects of polyamines on the reactions catalysed by, associated with rat ventral-prostate-gland chromatin and non-histone proteins (Ahmed, K., Wilson, M. J., Goueli, S. A. & Williams-Ashman, H. G.) 739–750

Protein modulator, calcium ion-dependent, properties of, of the activities of cyclic nucleotide phosphodiesterases of mouse epidermis (Murray, A. W. & Rogers, A.) 727–732

Protein, non-histone, high-mobility-group-type, high-molecular-weight, oviduct, chicken, isolation of (Teng, C. S., Gallagher, K. & Teng, C. T.) 1003–1006

Protein, oestradiol-17β-receptor, binding of the complex of oestradiol-17β with, to hypothalamus chromatin from male and female sheep (Perry, B. N. & Lopez, A.) 873–883

Protein, progesterone-receptor, binding of the complex of progesterone with, to hypothalamus chromatin from male and female sheep (Perry, B. N. & Lopez, A.) 873–883

Protein, progesterone-receptor, cytosolic, uterus, human, interaction of progestins with (Kasid, A., Buckshee, K., Hingorani, V. & Laumas, K. R.) 531–539

Protein, purification and properties of a factor inhibitory towards the biosynthesis of, from skeletal muscle of normal mice and those with muscular dystrophy (Petryshyn, R. A. & Nicholls, D. M.) 907–917

Protein, rates of biosynthesis of, in chicken skeletal muscle, heart muscle and smooth muscle (Laurent, G. J., Sparrow, M. P., Bates, P. C. & Millward, D. J.) 393–401

Protein, turnover of, in chick skeletal muscle during growth (Maruyama, K., Sunde, M. L. & Swick, R. W.) 573–582

Proteinases, sensitivity towards, of the release of α-amylase and ribonuclease by rat pancreatic microsomal fraction (Pearce, P. D., May, B. K. & Elliott, W. H.) 611–614

Protein–polysaccharides, *see* Proteoglycans

Proteins, collagenous, biosynthesis and partial characterization of, of rat lens capsule (Heathcote, J. G., Sear, C. H. J. & Grant, M. E.) 283–294

Proteins L7 and L12, ribosomal, *Escherichia coli*, characterization of the acidic ribosomal phosphoprotein L_7 from Krebs II ascites cells and its relationship to (Leader, D. P. & Coia, A. A.) 569–572

Proteins, membrane, erythrocyte, human, characterization of, after cell fusion induced by oleoylglycerol in the presence of calcium ions (Quirk, S. J., Ahkong, Q. F., Botham, G. M., Vos, J. & Lucy, J. A.) 159–167

Proteins, membrane, lysosomal, liver, rat, evidence that most of, are exposed to the cytoplasm (Schneider, D. L., Burnside, J., Gorga, F. R. & Nettleton, C. J.) 75–82

Proteins, mitochondrial, liver, rat, choice of precursors for the measurement of the turnover of proteins by a double-isotope method and its application to the study of (Burgess, R. J., Walker, J. H. & Mayer, R. J.) 919–926

Proteins, mitochondrial, liver, rat, relative rates of turnover of subunits of (Walker, J. H., Burgess, R. J. & Mayer, R. J.) 927–932

Proteins, non-histone, effects of polyamines on the reactions catalysed by protein kinases associated with chromatin and, of rat ventral-prostate-gland nuclei (Ahmed, K., Wilson, M. J., Goueli, S. A. & Williams-Ashman, H. G.) 739–750

Proteins, oestrogen-receptor, uterus, rat, binding characteristics, content and intracellular distribution of, in relation to intra-uterine devices and the oestrous cycle (Myatt, L., Chaudhuri, G., Elder, M. G. & Lim, L.) 523–529

Proteins, ribosomal, effect of various methods of preparation on the apparent composition of, of rabbit reticulocyte ribosomes (Huvos, P., Fey, S. & Hardwicke, P. M. D.) 381–391

Proteoglycans, cartilage, articular, human, age-related changes in the composition and structure of (Bayliss, M. T. & Ali, S. Y.) 683–693
Proteoglycans, sulphated, endocytosis of, by cultured human skin fibroblasts (Prinz, R., Schwermann, J., Buddecke, E. & von Figura, K.) 671–676
Protons, anion/calcium ion ratios and production of, during the uptake of calcium ions by rat heart and liver mitochondria (Harris, E. J.) 983–991
Pseudomonic acid, inhibition by, of the activity of isoleucyl-transfer ribonucleic acid synthetase in *Escherichia coli* M.R.E. 600 (Hughes, J. & Mellows, G.) 305–318
Pulmonary surfactant, accumulation and turnover of, in rabbit lung (Desai, R., Tetley, T. D., Curtis, C. G., Powell, G. M. & Richards, R. J.) 455–462
Purine nucleotides, systematic variations in the concentrations of, in steady-state perfused rat heart (Bates, D. J., Perrett, D. & Mowbray, J.) 485–493
Pyruvate dehydrogenase, stimulation by vasopressin of the activity of, in perfused rat liver (Hems, D. A., McCormack, J. G. & Denton, R. M.) 627–629
Pyruvate, exogenous, effect of, on the induction by glucose of the release of insulin by isolated rat islets of Langerhans (Sener, A., Kawazu, S., Hutton, J. C., Boschero, A. C., Devis, G., Somers, G., Herchuelz, A. & Malaisse, W. J.) 217–232
Pyruvate, stimulation by lactate and, of the biosynthesis of proinsulin and the release of insulin by isolated rat islets of Langerhans (Jain, K., Asina, S. & Logothetopoulos, J.) 31–37

Rana temporaria, *see* Frog
Red blood cells, *see* Erythrocytes
Redox state, effect of, of the cytosolic nicotinamide–adenine dinucleotide couple on the activity of adenosine 3′:5′-cyclic monophosphate phosphodiesterase and its role in the control of the responsiveness of isolated rat hepatocytes to glucagon (Clark, M. G. & Jarrett, I. G.) 805–816
Renin, interconversion of, and high-molecular-weight renin in dog kidney cortex (Funakawa, S., Funae, Y. & Yamamoto, K.) 977–981
Respiration, protective role of carotenoids in the photosensitivity of, by *Neurospora crassa* mitochondria (Ramadan-Talib, Z. & Prebble, J.) 767–775
Respiratory state, influence of, on the activity of monoamine oxidase in isolated rat liver mitochondria (Smith, G. S. & Reid, R. A.) 1011–1014
Reticulocytes, rabbit, effect of various methods of preparation on the apparent composition of ribosomal proteins of ribosomes from (Huvos, P., Fey, S. & Hardwicke, P. M. D.) 381–391
Reticulocytes, rat and rabbit, accelerated degradation of the protein formed in, under conditions of depletion of individual amino acids (Chandler, C. S. & Ballard, F. J.) 151–158
Reticulum, endoplasmic, liver, rat, evidence for the presence of a subpopulation of ribosomes bound to, that are detached *in vitro* by carcinogens and by centrifugation (Palmer, D. N., Rabin, B. R. & Williams, D. J.) 9–14
Ribonuclease, sensitivity towards proteinases of the release of α-amylase and, by rat pancreatic microsomal fraction (Pearce, P. D., May, B. K. & Elliott, W. H.) 611–614
Ribonucleic acid, effects of starvation on the turnover of protein and the concentration of, in active and immobilized rat skeletal muscle (Goldspink, D. F.) 603–606
Ribonucleic acid, high-molecular-weight, metabolism of, in hypothalamic and cortical regions of female rat brain during development (Hall, C. & Lim, L.) 511–521
Ribonucleic acid, initiation of the transcription of deoxyribonucleic acid for the biosynthesis of, in normal and phosphorylated rat liver nucleosomes (Ord, M. G. & Stocken, L. A.) 615–618
Ribonucleic acid, nuclear, heterogeneous, use of rat liver nucleoplasm for the characterization of the biosynthesis of, in rat liver *in vivo* (Beebee, T. J. C.) 715–725
Ribonucleic acid polymerase II, use of rat liver nucleoplasm for the characterization of the activity of, in rat liver *in vivo* (Beebee, T. J. C.) 715–725
Ribonucleic acid, ribosomal, onset of the degradation of, during cessation of growth in baby-hamster kidney BHK-21/C13 cells (Melvin, W. T. & Keir, H. M.) 933–941
[1-^{14}C]Ribose 5-phosphate, distribution of ^{14}C in glucose 6-phosphate biosynthesized from, by a rat liver extract and its relevance to the pentose phosphate pathway for the metabolism of glucose (Williams, J. F., Clark, M. G. & Blackmore, P. F.) 241–256
Ribosomal proteins L7 and L12, *Escherichia coli*, characterization of the acidic ribosomal phosphoprotein L_7 from Krebs II ascites cells and its relationship to (Leader, D. P. & Coia, A. A.) 569–572
Ribosomal proteins, *see also* Proteins, ribosomal
Ribosomal ribonucleic acid, *see* Ribonucleic acid, ribosomal
Ribosomes, brine-shrimp, inhibitory effect of modeccin on the elongation of polypeptide chains during the biosynthesis of protein catalysed by (Montanaro, L., Sperti, S., Zamboni, M., Denaro, M., Testoni, G., Gasperi-Campani, A. & Stirpe, F.) 371–379
Ribosomes, membrane-bound, liver, rat, evidence for the presence of a subpopulation of, that are detached *in vitro* by carcinogens and by centrifugation (Palmer, D. N., Rabin, B. R. & Williams, D. J.) 9–14
Ribosomes, onset of the degradation of, during cessation of growth in baby-hamster kidney BHK-21/C13 cells (Melvin, W. T. & Keir, H. M.) 933–941
Ribosomes, reticulocyte, rabbit, effect of various methods of preparation on the apparent composition of ribosomal proteins of (Huvos, P., Fey, S. & Hardwicke, P. M. D.) 381–391
Root nodules, kidney-bean, oxidation of glycine by leghaemoglobin from, and associated catabolic reactions at the haem group (Lehtovaara, P.) 351–358
Root nodules, leguminous-plant, formation of isomers of biliverdin during the degradation of leghaemoglobins from, of various species (Lehtovaara, P. & Perttilä, U.) 359–364

Salivary gland, parotid, rat, control of the biosynthesis and release of amylase by (McPherson, M. A. & Hales, C. N.) 855–863

Salivary gland, parotid, rat, dependence on calcium ions of the inhibition by α-adrenergic agonists and cholinergic agonists of the biosynthesis of protein in slices of (Kanagasuntheram, P. & Lim, S. C.) 23–29

Sarcoplasm, skeletal-muscle, rat, effect of surgical trauma on the turnover of protein in, and myofibrils (Hoover-Plow, J. L. & Clifford, A. J.) 137–142

[1-Sarcosine,8-alanine]angiotensin II, *see* Angiotensin II, [1-sarcosine,8-alanine]-

Serine, characteristics of the transport of alanine, glutamine and, across the plasma membrane of isolated rat hepatocytes (Joseph, S. K., Bradford, N. M. & McGivan, J. D.) 827–836

Serotonin, *see* 5-Hydroxytryptamine

Sex, variation of the concentrations of carnitine in various rat tissues with age and (Borum, P. R.) 677–681

Skeletal muscle, *see* Muscle, skeletal

Skin fibroblasts, human, cultured, endocytosis of sulphated proteoglycans by (Prinz, R., Schwermann, J., Buddecke, E. & von Figura, K.) 671–676

Skin fibroblasts, human, cultured, extensive conjugation of metabolites of 3,4-dihydroxyphenethylamine in, and rat hepatoma cells (Crooks, P. A., Breakefield, X. O., Sulens, C. H., Castiglione, C. M. & Coward, J. K.) 187–196

Skin, mouse, properties of the calcium ion-dependent protein modulator of the activities of cyclic nucleotide phosphodiesterases of (Murray. A. W. & Rogers, A.) 727–732

Small intestine, *see* Intestine, small

Smooth muscle, *see* Muscle, smooth

Somatotropin, stimulation by phospholipase A of the content of prostaglandin E and the biosynthesis of, in rat anterior pituitary gland (Betteridge, A. & Wallis, M.) 319–323

Soya bean, *see* Bean, soya

Spermidine, effect of, and other polyamines on the reactions catalysed by protein kinases associated with rat ventral-prostate-gland chromatin and non-histone proteins (Ahmed, K., Wilson, M. J., Goueli, S. A. & Williams-Ashman, H. G.) 739–750

Spermidine, pathway for the biosynthesis of, and other polyamines in *Caldariella acidophila* (De Rosa, M., De Rosa, S., Gambacorta, A., Cartenì-Farina, M. & Zappia, V.) 1–7

Spermine, effect of, and other polyamines on the reactions catalysed by protein kinases associated with rat ventral-prostate-gland chromatin and non-histone proteins (Ahmed, K., Wilson, M. J., Goueli, S. A. & Williams-Ashman, H. G.) 739–750

Sporulation, accumulation of calcium ions during, of *Bacillus megaterium* KM (Hogarth, C. & Ellar, D. J.) 197–203

Starvation, effects of, on the turnover of protein and the concentration of ribonucleic acid in active and immobilized rat skeletal muscle (Goldspink, D. F.) 603–606

Starvation, opposite effects of, on the oxidation of adenosine and on adenosine-induced release of insulin by isolated mouse islets of Langerhans (Andersson, A.) 619–621

Steroids, effects of angiotensin II and [1-sarcosine,8-alanine]angiotensin II on the production of adenosine 3′:5′-cyclic monophosphate and the biosynthesis of, by isolated rat adrenal-gland-cortex glomerulosa cells (Bing, R. F. & Schulster, D.) 39–45

Stomach, human, digestion of ovalbumin by pepsins 1, 3 and 5 from (Walker, V. & Taylor, W. H.) 429–432

Stomach, rat, possible role of 5-hydroxytryptamine as mediator in the action of calcitonin on, and small intestine and pancreas (Nakhla, A. M. & Latif, A.) 339–342

Streptozotocin-diabetes, *see* Diabetes, streptozotocin-induced

Sulphate, inorganic, activity of phenol sulphotransferase and the metabolism of, in isolated perfused rat liver (Mulder, G. J. & Keulemans, K.) 959–965

Supernatant fraction, *see* Cytosol

Synaptosomes, brain, guinea-pig, activation by calcium ions of the hydrolysis of phosphatidyl-*myo*-inositol 4-phosphate (diphosphoinoside) and phosphatidyl-*myo*-inositol 4,5-bisphosphate (triphosphoinositide) by (Griffin, H. D. & Hawthorne, J. N.) 541–552

Synaptosomes, brain, rat, rapid method for the isolation of a relatively pure metabolically competent preparation of (Booth, R. F. G. & Clark, J. B.) 365–370

Testis, mouse, induction by the β-subunit of human chorionic gonadotropin of the production of testosterone by Leydig cells from (Ramakrishnan, S., Das, C. & Talwar, G. P.) 599–602

Testosterone, induction by the β-subunit of human chorionic gonadotropin of the production of, by mouse testis Leydig cells (Ramakrishnan, S., Das, C. & Talwar, G. P.) 599–602

Thromboxane synthase, evidence that the formation of malondialdehyde in human platelets is catalysed by (McMillan, R. M., MacIntyre, D. E., Booth, A. & Gordon, J. L.) 595–598

Thyroid hormones, modulation by, of the numbers of β-adrenergic receptors in membranes of isolated rat epididymal adipocytes (Giudicelli, Y.) 1007–1010

Thyroxine, effects of cortisol, corticotropin and, on the biosynthesis of glycerolipids and on the activity of phosphatidate phosphatase in rat liver (Glenny, H. P. & Brindley, D. N.) 777–784

Thyroxine, effects of, on the activity of phosphatidate phosphatase in rat liver after the administration of ethanol and glycerol to the animals (Savolainen, M. J. & Hassinen, I. E.) 885–892

Toad (*Bufo marinus*), effects of cholera toxin and guanosine 5′-[βγ-imido]triphosphate on the affinity of the β-adrenergic receptors of erythrocyte membranes from (Fischer, J. & Sharp, G. W. G.) 505–510

Trauma, surgical, effect of, on the turnover of protein in rat skeletal muscle (Hoover-Plow, J. L. & Clifford, A. J.) 137–142

Triacylglycerols, inactivation of rat epididymal adipose-tissue glycerol phosphate acyltransferase catalysed by adenosine 3′:5′-cyclic monophosphate-dependent protein kinase and its possible involvement in the regulation of the biosynthesis of (Nimmo, H. G. & Houston, B.) 607–610

Triacylglycerols, stimulation by human very-low-density plasma lipoproteins of the biosynthesis of, in isolated rat adipocytes (Thomopoulos, P., Berthelier, M., Lagrange, D., Chapman, M. J. & Laudat, M.-H.) 169–174
Triglycerides, *see* Triacylglycerols
3,3′,5′-Tri-iodothyronine, modulation by, of the numbers of β-adrenergic receptors in membranes of isolated rat epididymal adipocytes (Giudicelli, Y.) 1007–1010
Triphosphoinositide, *see* Phosphatidyl-*myo*-inositol 4,5-bisphosphate
Tryptophan, differential effects of, on gluconeogenesis in isolated rat and guinea-pig hepatocytes (Smith, S. A., Elliott, K. R. F. & Pogson, C. I.) 817–825
Tyrosine aminotransferase, participation of cysteine and cystine in the inactivation of, in rat liver homogenates (Buckley, W. T. & Milligan, L. P.) 449–454

Ubiquinone-8, deficiency in the pathway for the biosynthesis of, in a mutant of *Escherichia coli* K12 that is sensitive to chlorate when grown aerobically (Giordano, G., Grillet, L., Rosset, R., Dou, J. H., Azoulay, E. & Haddock, B. A.) 553–561
Urea, effects of alloxan-induced diabetes on the formation of glucose and, in isolated rabbit liver cells (Zaleski, J. & Bryła, J.) 563–568
Urea, sources of ammonia for the biosynthesis of, by isolated rat hepatocytes (Krebs, H. A., Hems, R., Lund, P., Halliday, D. & Read, W. W. C.) 733–737
Ureogenesis, sources of ammonia for, by isolated rat hepatocytes (Krebs, H. A., Hems, R., Lund, P., Halliday, D. & Read, W. W. C.) 733–737
Uterus, human, interaction of progestins with the cytosolic progesterone-receptor protein of (Kasid, A., Buckshee, K., Hingorani, V. & Laumas, K. R.) 531–539
Uterus, rat, binding characteristics, content and intracellular distribution of oestrogen-receptor proteins in, in relation to intra-uterine devices and the oestrous cycle (Myatt, L., Chaudhuri, G., Elder, M. G. & Lim, L.) 523–529

Valine, accelerated degradation of protein formed in rat and rabbit reticulocytes under conditions of depletion of, and other individual amino acids (Chandler, C. S. & Ballard, F. J.) 151–158
Vasopressin, effects of phenylephrine, glucagon and, on gluconeogenesis and on enzymes involved in the metabolism of glycogen in isolated rat hepatocytes (Hue, L., Felíu, J. E. & Hers, H.-G.) 791–797
Vasopressin, stimulation by angiotensin, oxytocin and, of gluconeogenesis in isolated rat hepatocytes (Whitton, P. D., Rodrigues, L. M. & Hems, D. A.) 893–898
Vasopressin, stimulation by, of the activity of pyruvate dehydrogenase in perfused rat liver (Hems, D. A., McCormack, J. G. & Denton, R. M.) 627–629
Vicia faba, *see* Bean, broad
Vitamin D_3, *see* Cholecalciferol